

Original research article

A Study on Knowledge, Attitude and Perceptions of 1st year MBBS Students about Hepatitis B Virus Infection and Vaccination

¹Dr. Arifuddin MK , ² Dr.Mir Viqar Ahmed

¹Assistant Professor, Department of Physiology, Sri Siddhartha Medical College, Tumkur, India

²Assistant Professor, Department of Community Medicine, Sri Siddhartha Medical College, Tumkur, Karnataka, India

Corresponding Author: Dr. Arifuddin MK

Abstract

Background: Hepatitis B virus infection is one of the important causes for the development of chronic liver disease. The prevalence of chronic Hep- B virus infection in India is 1.46% with an estimated 17 million people who are chronic carriers of Hepatitis virus. Compared to the general public, the Seroprevalence of HBV infection is quiet higher among health care providers.

Objectives: To assess the knowledge,attitude and perceptions of 1st year MBBS students regarding HBV infection and vaccination.

Methodology: This study was carried out among 1st MBBS students belonging to 2015-16 batch of SSMC, Tumkur, Karnataka. A self-structured objective type of questionnaire was prepared consisting of questions which were used to assess the knowledge regarding HBV infection and vaccination for it. The responses were analyzed using Microsoft excel and Epi Info 7.

Results: 92.3% of the students told that they have heard about hepatitis and 78.46% were also aware that the primary organ affected in hepatitis is liver. 87.69% responded that they were aware of hepatitis being transmitted from person to person. 89.23% of the study population had correct knowledge about the mode of transmission.

Conclusion: It is recommended to increase the awareness of importance of vaccination against HBV at the beginning of the medical course and consider to vaccinate all of them before they are exposed to the patients during their clinical postings in near future.

Keywords: Hepatitis B Virus infection, chronic liver disease, medical students, vaccination

Introduction

Hepatitis B virus infection which is one of the major causes of chronic liver disease and it still continues to assume endemic proportions in many countries of the world inspite of implementation of vaccination program for the same. The World Health Assembly has called for a goal for eliminating viral hepatitis as a major public health burden by the year 2030 [1]. It is estimated that 257 million people are living with chronic hepatitis B infection globally and 8,07,000 deaths were estimated due to complications arising from chronic hepatitis B infection in the year 2015 [2]. The prevalence of chronic hepatitis B virus infection in India is 1.46% with an estimated 17 million people who are chronic carriers of Hepatitis viral infection [3]. WHO recommends vaccination for hepatitis B virus at the birth which is followed by 2 or 3 doses [4]. Studies done in many other countries have shown a marked reduction in the prevalence of chronic HB virus infection and the occurrence of hepatocellular carcinoma in fully vaccinated people [5, 6]. Seroprevalence of HBV infection is few times higher among health care workers when compared to the general population [7]. The high prevalence of HBV infection is attributed to issues like incorrect handling of infectious materials and issues regarding improper sterilization of medical equipments [8, 9]. Medical students and house surgeons are at a greater risk of contracting HBV infection because of their inexperience in handling the infected material [10, 11]. Hepatitis B vaccination program was launched in 2003 in India and it was expanded to the entire country in 2008 [12]. It is unlikely that the Indian population born before 2008 have not received Hepatitis B vaccination as it was not a part of Immunization program. Several studies done earlier have indicated that there is a clear gap of knowledge and information about safe medical practices to reduce the risks of occupational exposure to HBV infection [13, 14]. Hence the present study is taken up among the first year medical students to assess the knowledge and their perceptions regarding HBV infection and vaccination for the same.

MATERIALS AND METHODS

This cross sectional study was carried out among first year MBBS students belonging to 2016-17 batches of Sri Siddhartha Medical College, Tumkur, and Karnataka after obtaining the approval of institutional ethical committee. All the 130 students were enrolled in the study. A self-structured objective type of questionnaire was prepared consisting of questions which were used to assess the knowledge regarding HBV infection and vaccination for it. Informed consent was taken and the confidentiality was maintained during the process of this survey. The responses were analyzed using Microsoft excel and Epi Info 7.

RESULTS

Table 1: Age and gender distribution

	Mean	Standard deviation
Age of the subject	18.05	0.52
Sex	Frequency	Percentage
Female	72	55.38
Male	58	44.61

Table 2: Respondents knowledge about HBV infection and vaccination

Have you heard about hepatitis?	120 (92.3%)
What organs is primarily affected in hepatitis-B?	102 (78.46%)
Can hepatitis-B transmitted from one person to another?	114 (87.69%)
What in your view is / are the mode of its transmission? (sexual & parental)	116 (89.23%)
Who are at higher risk of acquiring hepatitis	78 (60%)
Is hepatitis-B infection preventable by vaccination	98 (75.38%)

DISCUSSION

Hepatitis B is an acute viral infection which is caused by HBV primarily affecting the liver with potential complications which include hepatocellular carcinoma, cirrhosis of liver and chronic hepatitis [15,16]. Healthcare professionals are at an increased risk of occupational exposure to HBV infection especially in developing countries. Our study population included 130 first year medical students belonging to 2016-17 batch. The average age distribution of the study population was 18.05 years and males constituted 58 (44.61%) and females 72 (55.38%) of the study population. Analysis of the responses revealed that 92.3% of the study population told that they have heard about hepatitis and 78.46% were also aware that the primary organ affected in hepatitis was liver. 87.69% responded that they were aware of hepatitis being transmitted from person to person. 89.23% of the study population had correct knowledge about the mode of transmission. These findings are consistent with another study done by Darwish and AL Khaldi [17] done among medical students. In another study done by Sanjay Kumar Choudhary and et al. revealed that 92% of the students had correct knowledge regarding the mode of transmission [18]. 60% of the study population was aware of the higher risk of acquiring HBV infection among healthcare workers. 75.38% of the students reported that they were aware that HBV infection can be prevented by vaccination although only 46% of the study population was aware of the correct Hepatitis B immunization schedule. This is in consistent with another study done by Sanjay Kumar Choudhary and et al. where 52% of the study population did not know about the correct vaccination schedule [18].

CONCLUSION

HBV infection which primarily affects the liver with a potential of causing chronic diseases can be effectively prevented by vaccination. Our study provides many evidences about the knowledge and perceptions about HBV infection in first MBBS students. In spite of having a good knowledge about the infection and the mode of transmission and also with the availability of vaccine it was found that 69.23% of students were not immunized. It is recommended to increase the awareness of importance of vaccination against HBV at the beginning of the medical course and consider vaccinating all of them before they are exposed to the patients during their clinical postings.

REFERENCES

1. World Health Organization. (2017). WHO Global Hepatitis Report. Geneva: World Health Organization.
2. World Health Organization. (2017). World Hepatitis Report. 2017 Geneva. Available from: <https://apps.who.int/iris/bitstream/handle/10665/255016/9789241565455eng.pdf;jsessionid=10847CB8AB36884692B8BEBBD0D54EB5?sequence=1>, accessed Aug 17, 2020.
3. Schweitzer, A., Horn, J., Mikolajczyk, R. T., Krause, G., & Ott, J. J. (2015). Estimations of worldwide prevalence of chronic hepatitis B virus infection: a systematic review of data published between 1965 and 2013. *The Lancet*, 386(10003), 1546-1555.
4. World Health Organization. (2019). Hepatitis B vaccines: WHO position paper, July 2017–Recommendations. *Vaccine*, 37(2), 223-225.
5. Chang, M. H., Chen, C. J., Lai, M. S., Hsu, H. M., Wu, T. C., Kong, M. S., ... & Chen, D. S. (1997). Universal hepatitis B vaccination in Taiwan and the incidence of hepatocellular carcinoma in children. *New England Journal of Medicine*, 336(26), 1855-1859.
6. Liang, X., Bi, S., Yang, W., Wang, L., Cui, G., Cui, F., ... & Wang, Y. (2013). Reprint of: Epidemiological serosurvey of Hepatitis B in China—declining HBV prevalence due to Hepatitis B vaccination. *Vaccine*, 31, J21-J28.
7. Singhal, V., Bora, D., & Singh, S. (2009). Hepatitis B in health care workers: Indian scenario. *Journal of Laboratory physicians*, 1(2), 41.
8. Prüss-Üstün, A., Rapiti, E., & Hutin, Y. (2005). Estimation of the global burden of disease attributable to contaminated sharps injuries among health-care workers. *American journal of industrial medicine*, 48(6), 482-490.
9. Taneja, N., & Biswal, M. (2006). Safe disposal of infectious waste—Indian perspective. *Journal of Hospital Infection*, 62(4), 525-527.

10. Rogers, B., & Goodno, L. (2000). Evaluation of interventions to prevent needlestick injuries in health care occupations. *American Journal of Preventive Medicine*, 18(4), 90-98.
11. Kohn, W. G., Collins, A. S., Cleveland, J. L., Harte, J. A., Eklund, K. J., & Malvitz, D. M. (2003). *Guidelines for infection control in dental health-care settings-2003*.
12. Lahariya, C., Subramanya, B. P., & Sosler, S. (2013). An assessment of hepatitis B vaccine introduction in India: Lessons for roll out and scale up of new vaccines in immunization programs. *Indian journal of public health*, 57(1), 8.
13. Pathoumthong, K., Khampanisong, P., Quet, F., Latthaphasavang, V., Souvong, V., & Buisson, Y. (2014). Vaccination status, knowledge and awareness towards hepatitis B among students of health professions in Vientiane, Lao PDR. *Vaccine*, 32(39), 4993-4999.
14. Noubiap, J. J. N., Nansseu, J. R. N., Kengne, K. K., Ndoula, S. T., & Agyingi, L. A. (2013). Occupational exposure to blood, hepatitis B vaccine knowledge and uptake among medical students in Cameroon. *BMC medical education*, 13(1), 1-5.
15. Barker, L. F., Shulman, N. R., Murray, R., Hirschman, R. J., Ratner, F., Diefenbach, W. C., & Geller, H. M. (1970). Transmission of serum hepatitis. *Jama*, 211(9), 1509-1512.
16. Ozaras, R., & Tahan, V. (2009). Acute hepatitis C: prevention and treatment. *Expert review of anti-infective therapy*, 7(3), 351-361.
17. Darwish, M. A., & Al Khaldi, N. M. (2013). Knowledge about hepatitis B virus infection among medical students in university of dammam, eastern region of Saudi Arabia. *Life Sci J*, 10(2), 860-7.
18. Choudhary, S. K., Kumar, S., & Sinha, S. (2017). Knowledge and Vaccination Coverage of Hepatitis-B among 1st Year MBBS Students at Indira Gandhi Institute of Medical Sciences, Patna. *International journal of scientific study*, 4(10), 101-104.