

# Management of distal tibial intra-articular fractures by using ring external fixators assisted arthroscopically

Mohammed Safwat Shalabi, Mohammed Abdel Wahab Ibrahim, Ashraf Abd Al-Daim  
Mohamed, Mohammed Osama Mohammed Morsi\*

Departments of Orthopaedic Surgery, Faculty of Medicine - Zagazig University

\*Corresponding author: Mohammed Osama Mohammed Morsi,

Mobile: (+20) 0107682955, E-Mail: amanyohed77@gmail.com

*Background: Tibial pilon fractures are relatively uncommon injuries, representing only 1% of all fractures of the lower limb and 5% to 10% of those of the tibia. Frequent comminution and the thin soft-tissue envelope in the area make the treatment of these fractures challenging. The tibial pilon is characterised by a total absence of muscle coverage and marginal vascularity, therefore, even moderate trauma often results in extensive soft-tissue damage.*

*Objective: The aim of this present study was to evaluate the arthroscopic assisted (Ilizarov) ring external fixation of distal tibial intra articular (pilon) fractures.*

*Patients and methods: This was a prospective study conducted between February 2012 and April 2015 on thirty patients with closed and Gustilo Types I & II open fractures of pilon fractures of the distal tibia who were admitted to Zagazig University Hospitals. during a period of two years and all cases were treated by Ilizarov fixators with or without limited internal fixation and assessment of intra-articular reduction tibial plafond by arthroscopy.*

*Results: Nineteen patients had right-sided injury, eleven patients had left sided injury and one patient had bilateral injury. At the time of injury the youngest patient was 19 years old and the oldest was 60 years. The results in relation to pain as distributed according to the fracture, it is noted that the excellent results were with type I, II fracture while the fair and poor results were with type II and III fracture.*

*Conclusion: The present study provides information on the use and the role of ankle arthroscopy in assisting and evaluation of distal tibial intra articular reduction after pilon fracture, also the value, advantages, techniques of ankle arthroscopy and ring external fixators. The study showed that this technique might be a good method to manage the pilon fractures.*

*Keywords: Distal tibial intra-articular fractures, Pilon fractures, Arthroscopically*

## Introduction

Two distinct mechanisms are believed to be responsible for most tibial pilon fractures: low-stress trauma (sports injuries), which is less common and secondary to rotational forces, and high-stress trauma (motor vehicle accidents, falls from height, workplace accidents), which is more common and produces axial transmission of the load with the talus pushed onto the distal tibia, resulting in a multi fragmentary implosion of bones and cartilage structures<sup>(1)</sup>. The pattern of the fracture with this mechanism of injury depends on the position of the foot at the time of trauma. With the foot dorsiflexed, compressive forces are placed on the anterior part of the plafond, whereas with the foot plantarflexed the forces are directed

posteriorly. If the foot is in a neutral position, the axial force can also compromise the integrity of the articular surface <sup>(1)</sup>.

Several classification systems have been proposed for tibial pilon fractures. Rüedi and Allgöwer introduced a classification that divides fractures into three groups according to the degree of displacement and comminution of the articular and metaphyseal fragments <sup>(2)</sup>, to which Ovadia and Beals added two fracture variants <sup>(3)</sup>. Currently, the most widely used classification is the AO/OTA classification, which divides fractures into three groups: type A, extra-articular; type B, partially intra-articular; and type C, completely intra-articular. Each group is further divided into subgroups that indicate the severity of comminution and the fracture pattern <sup>(4)</sup>.

The treatment of tibial pilon fractures, particularly high-stress fractures, is complex and often difficult. The historically disappointing results of conservative treatment and the first attempts at limited internal fixation led the AO to develop guidelines for open reduction and internal fixation. This method has produced and still produces good results, particularly in less severe, low stress fractures, but results are not as good in high-stress injuries <sup>(5)</sup>.

In fractures with severe soft-tissue damage and metaphyseal comminution, the introduction of plates and screws through an extensive exposure is associated with a high rate of failure and severe complications, such as surgical wound infections, osteomyelitis, and non-union <sup>(5)</sup>.

Hybrid external fixation represent a possible alternative method to provide stable fixation and reduce complications. These devices have gained increasing popularity among orthopaedic surgeons because they provide a stable fracture synthesis that can drastically reduce soft-tissue damage; however, in some cases, a relatively simple and standardized method of assembly of the external fixation is balanced by the need for minimal additional internal fixation to restore proper joint congruity. Despite the growing interest in this type of technique, most of the studies in the literature are based on relatively small numbers of cases <sup>(6)</sup>.

The aim of this present study was to evaluate the arthroscopic assisted (Ilizarov) ring external fixation of distal tibial intra articular (pilon) fractures. Using the following approach: reduction of the fracture, visualization of the plafond articular fracture by standard ankle arthroscopic techniques, and application of ring external fixators.

### **Patients and methods**

This was a prospective study conducted between February 2012 and April 2015 on Thirty patients with closed and Gustilo Types I&II open fractures of pilon fractures of the distal tibia who were admitted to Zagazig University Hospitals. during a period of two years and all cases were treated by Ilizarov fixators with or without limited internal fixation and assessment of intra-articular reduction tibial plafond by arthroscopy. And were followed up for one year with an average 6 months, the majority occurring in males. The average age was 35(sd±11) years and range from 19 years to 60 years. The data of each patient were collected as regard the name, age, Sex, occupation, address, special habits, mechanism of injury, associated injury, history of receiving treatment of diabetes, hypertension, and peripheral vascular disease.

**Diagnosis**

In diagnosis of our patients complete history and physical examination was taken. The specific details of the mechanism of injury are important as they will suggest the type of force exerted upon both the soft and skeletal tissues which will vary considerably depending upon the mechanism of injury.

**History:**

- a) Age of the patient
- b) Mechanism of injury
  - The history of jumping injury or an accident or a simple falls at home.
  - Information that is relevant to the mechanism of injuries includes: Distance of the fall, surface onto which the patient fall and the position in which the patient landed.

Sever compression injuries were elicited in patients falling from height (four cases), or Motor vehicle-related accidents (six cases). Minimal comminution and large articular fragments were seen in with rotational injuries like falling from height (low distance) (14cases), fall down stairs (three cases) and falling forward with entrapped foot (three cases).

Some systemic disease which may be associated with vascular complication such as diabetes mellitus or systemic vascular disease. We have: Two diabetic patients one of them have hypertension also. Three patients have hypertension.

**Clinical examination:**

- Patients had been examined for any associated injuries.
- Clinical examination first assess the condition of the skin (closed or open)

**Swelling:**

Swelling may be produced by direct acting external force and by internal fragment pressure or fracture haematoma. In pilon swelling is collar like imbibitions of the distal lower leg contrary to malleolar fracture where the swelling presents as a haemoarthrosis, so the patients around the ankle joint. Swelling and ecchymosis under the planter aspect of the foot was found in a case with fracture calcaneus. Ten cases with skin contusion and abrasion, nine cases with skin blisters.

**Vascular assessment:**

Our examinations also include evaluation of the peripheral arterial on as the swelling may be precluding palpation the foot pulse our based on the microcirculation (toes). There are no cases with impairment or injury of major blood vessels. Motor function tested by active movement of the toes particularly the ankle, metatarsus and calcaneus are too close to the pain radiating from fracture site so difficult to test.

**Neurological status:**

Includes both planter and dorsal aspect of the foot sensorimotor status of the foot due to fracture was not found in our cases. But we had two cases with bilateral hypothesia due to diabetes.

**Roentgenogram examination:**

Pain x-ray film of the affected leg includes the entire tibia and fibula for all patients. Antero-posterior, lateral and multiple oblique views for all the patients. Plain x-ray film for unaffected leg was taken.

Roentgenogram had been examined to identify:

- i. Fracture line, the major fracture fragment and the number of fragments.
- ii. Pieces of subchondral bone, which may have been driven into the assessment of bone quality.

**Statistical analysis:**

All data were collected, tabulated and statistically analyzed using SPSS 18.0 for windows (SPSS Inc., Chicago, IL, USA). Continuous data are expressed as the mean  $\pm$  SD & median (range), and the categorical data are expressed as a number (percentage). Continuous variables were checked for normality by using Shapiro-Wilk test. Independent samples Student's t-test was used to compare two groups of normally distributed data. Mann Whitney U test was used to compare two groups of non-normally distributed data. One way ANOVA test was used to compare more than two groups of normally distributed data. Kraskall Wallis H test was used to compare more than two groups of non-normally distributed data. Categorical data were compared using the Chi-square ( $\chi^2$ ) test or Fisher's exact test when appropriate. All tests were two tailed.  $p < 0.05$  was considered statistically significant (S),  $p < 0.01$  was considered highly statistically significant (HS), and  $p > 0.05$  was considered non statistically significant (NS).

**Results**

At the time of injury the youngest patient was 19 years old and the oldest was 60 years. Table (1) shows classification of patients according to their ages.

**Table (1): Age of the patients.**

Age in year	Number of patient	Percentage
18-19 y	2	7%
20-29 y	9	30%
30-39 y	10	33%
40- 49 y	6	20%
> 50 y	3	10%

**Table (2): Sex and side distribution.**

Sex	No.	Right	Left
Male	24	15	9
Female	6	4	2
<b>Total</b>	30	19	11

Table (2) shows gender of the patients and side of fractures. Nineteen patients had right-sided injury, eleven patients had left sided injury and one patient had bilateral injury.

**Table (3): Mechanism of injury.**

Cause of trauma	No.	Percentage
Road traffic accidents	4	13%
Fall from height	20	70%
Fall on stairs	3	10%
Fall with entrapped foot	3	7%

Table (3) shows the causes of injuries. Height of fall varied approximately from 2 meter to 6 meters Reported patients and their-relatives.

Associated injuries: table (4)

- Associated injuries were present in 12 patients (40%), some of them suffered from two or three sites of associated skeletal injuries (3 cases).
- Patient no. 6 associated with fracture distal femur plus wedge fracture of 1<sup>st</sup> lumbar Vertebra the cause was.
- Patient no. 11 associated with fracture of ipsilateral forearm and fracture of contra lateral tibia after car accident.
- Case no. 20 associated with contra lateral fractures tibial shaft and distal radius after falling from height.

**Table (4): Associated injuries.**

Associated injuries	No. Of the patients
Tibial plateau fracture	1
Vertebral fracture	2
Forearm fracture	1
Olecranon fracture	1
tibial shaft fracture	2
Bimalleolar fracture	1
Os calcis fracture	2
Colles' fracture	1
Fracture femur	2
Fracture neck-femur	1
Brain concussion	2

#### Classification:

The plain x-ray was studied and the fracture was classified according to **Rüedi and Allgöwer**<sup>(2)</sup> classification, which based on the degree of displacement of articular fragments.

- Two patients (6.67%) had type I fracture both of them are males.
- Nineteen patients (63-33%) had type II fracture, five of them are females and fourteen are males. (Table 5).

**Table (5): Classification.**

Classification of fracture	Number of patients	Percentage
Type I	2	6.6%
Type II	19	63.4%
Type III	9	30%

Table (5) shows types of fractures and number of the patients. Nine patients (30%) with type III, two patients are female and seven patients are males. (Table 6).

**Table (6): Number of male and female patients.**

Classification of fracture	Number of patients	Male	Female
Type I	2	2	-
Type II	19	14	5
Type III	9	7	2
<b>Total</b>	<b>30</b>	<b>23</b>	<b>7</b>

**Closed or open fractures:**

Number of patients had open fractures:

With type II pilon fracture, one with Gustilo type one open fracture, other with Gustilo type three A, open fracture. Third patient had (type) fracture and type one open fracture) and fourth patient had open fibula type two open fracture (type three pilon fracture). (Table 7)

**Table (7): Number of open and closed fractures.**

Classification of fracture	Number of patients	Male	Female
Type I	2	-	2
Type II	19	2	17
Type III	9	2	7
<b>Total</b>	<b>30</b>	<b>4</b>	<b>26</b>

Types I fractures were torsion injuries whereas type II and III were associated with vertical compression (fall from height, road traffic accident).

The results in relation to pain as distributed according to the fracture, it is noted that the excellent results were with type I, II fracture while the fair and poor results were with type II and III fracture (Table 8).

**Table (8): Results in relation to pain.**

	Number of the patients		Type of fracture		
	No.	Percent	I	II	III
<b>Excellent</b>	5	16.7%	2	3	
<b>Good</b>	17	56.7%	--	14	3
<b>Fair</b>	6	20.0%		2	4
<b>Poor</b>	2	6.6%			2

**Discussion**

Although open reduction and anatomical reconstruction of the articular surface is ideal in any intraarticular fracture, it is particularly difficult in high-energy tibial pilon fractures. Pre-existing severe soft tissue injury will preclude open reduction in view of the potential wound problems due to ischaemia or infection of the soft tissues. Precarious blood supply in this region of the tibia can add to the problems of nonunion along with infection <sup>(7)</sup>.

We believe that treatment with Ilizarov external fixator provides multiple advantages in regard to fragment manipulation, stabilization, and maintenance of capsular and ligamentous traction required to enhance and promote normal cartilage metabolism <sup>(8)</sup>.

In a recent study of 23 patients with high-energy tibial pilon fractures treated with early external ring fixation and arthroscopic assisted reduction, **EL-Mowafi et al.** <sup>(9)</sup> found encouraging results with good functional outcome, (as well as reduced rates of ankle stiffness) There were no long-term problems with fracture union, and no patient required an ankle fusion at the end of 1 year of follow-up. Compared with their patients our series had less severely injured patients and hence better outcomes. The Ilizarov percutaneous fixator, preserves endosteal and periosteal blood supply, helps capture the small metaphyseal and subchondral bony fragments, and also helps compression of fracture fragments using the olive wires. The rigidity of fixation can be adjusted to suit the stage of fracture healing. It also allows correction of deformity during the process of fracture healing.

Arthroscopic assisted fracture reduction can provide a less invasive operative reduction of the fracture and respects the soft tissue envelope. **Lonjon et al.** <sup>(10)</sup> in a techniques similar to ours, describe ankle arthroscopy as useful tool and seemed more appropriate to us to use this minimal surgery, we used a standard 4-mm arthroscope through the classic portal of anterior arthroscopy.

Most of our cases are males 83 %, with main age 36 years , this is explained by that lesion is work related. Regarding to the fracture type more than half of the patients have type II Ruedi fracture, the incidence of open fractures is 83% . **Zhao et al.** <sup>(11)</sup> in a retrospective study on Twenty-five type C pilon fractures, including 4 open fractures, using the same technique had results An excellent or good AOFAS outcome was obtained in 81% of the patients. most of the studies that deals with this type of fractures and use the arthroscopic assisted ring fixation lack the high percentage of open fractures in comparison to our study.

Multiple strategies have been employed for dealing with the challenges of open pilon fractures. **Sirkin et al.** <sup>(12)</sup> reported on a series of pilon fractures using plate or Ilizarov-type constructs However, they used the Ilizarov system exclusively in their open fractures, which the authors attributed to their low wound complication rate.

The main time passed before the operation, in our sries, is 3.5 days, with median range 3(1-7) due to the preparations and management of other associated lesions, with consideration of changing the operative procedure as some cases were planned to open reduction and ineternal fixation but edema and skin bullae were challenging, recently some articles have recommended early interference before edema occurs. In the study of **Kapoor et al.** <sup>(13)</sup> on 17 patients with type II and III pilon fracture the main interval between injury and operation was 3.4 (6 hours to 23 days), they emphasised that of time of surgical interference with ilizarov fixation is not so important as open reduction and fixation by plate, while **Sirkin et al.** <sup>(12)</sup> considered that thin wire ring external fixator (with or without internal fixator) is preffered to be used within the first 72 hours after injury.

In a recent study of 23 patients with high-energy tibial pilon fractures treated with early external ring fixation and arthroscopic assisted reduction, **EL-Mowafi et al.** <sup>(9)</sup> found encouraging results with good functional outcome, (as well as reduced rates of ankle stiffness)

Arthroscopic assisted fracture reduction can provide a less invasive operative reduction of the fracture and respects the soft tissue envelope. **Guillo et al.** <sup>(14)</sup> in a technique similar to ours, describe ankle arthroscopy as useful tool and seemed more appropriate to us to use this

minimal surgery, we used a standard 4-mm arthroscope through the classic portal of anterior arthroscopy.

The fragments are then fixed to their anatomical place by Kirschner wires and olive wires adjusted properly to the ring external fixator, as **Vidyadhara and Rao** <sup>(7)</sup> recommended this technique as olive wires squeeze the fracture pattern in a very methodical way, they treated 21 consecutive patients with complex tibial pilon fractures, using percutaneous reduction and fixation with Ilizarov apparatus.

In the present study, we depend on both ankle arthroscopy and fluoroscopy to help and evaluate the reduction, by direct vision and radiologically. Many studies were released considering the reduction of pilon fracture by ring external fixators, but yet very few studies were conducted to evaluate and describe the combined arthroscopic technique with external fixators in the treatment of distal tibial intra articular fractures, **EL-Mowafi et al.** <sup>(9)</sup>, enrolled 23 patients in their study, using the same technique.

**Lonjon et al.** <sup>(10)</sup> in their case presentation, showed that minimally invasive surgical reduction under arthroscopic guidance appears to be a promising alternative in this type of fracture.

In our study, the ankle syndesmosis, talar dome cartilage and deltoid ligament integrity were examined, in 21 patients (70%) these pathological entities were absent, while talar dome cartilage lesions (osteochondral flaps, ulcers) were found in 5 patients (16,7%) ankle syndesmosis was examined radiologically and arthroscopically to exclude occult diastasis, torn inter syndesmotric ligaments and anteroinferior tibiofibular ligaments. syndesmotric disruption was confined if the interval between tibia and fibula, according to **Chiang et al.** <sup>(15)</sup>, is increased by more than 2 mm when viewed from the anterolateral ankle portal.

**Chiang et al.** <sup>(15)</sup> described an algorithm for arthroscopic assisted reduction and evaluation of the distal intra articular tibial fractures, ankle syndesmosis and ankle ligaments especially the deltoid ligament, in their retrospective study on 105 patients with distal intra articular tibial fractures, divided into two groups, ankle arthroscopy was used as diagnostic tool for ankle structures evaluation in the first group only, the found in the first group that 52 patients (80%) had ankle syndesmotric distasis that need fixation, while the same lesion was discovered in only 23 (57.5%) patients in the second group.

Associated fibular fracture with ipsilateral pilon fracture, in our study, was present in 27 patients (90%), with slightly higher incidence compared with previous studies, as they conducted that the incidence is about 80% **Tomás-Hernández** <sup>(16)</sup> and **Busel et al.** <sup>(17)</sup>.

Tibial pilon fractures are frequently accompanied by a rotational component in addition to axial load, which should be resisted with a chosen fixation method. Rotational forces not only affect the distal tibia but often result in a fibular fracture as well, allowing for a marker during radiographic evaluation. comminuted fibular fractures are seen as a result of valgus stresses, whereas transverse fractures of the fibula result from varus angulation, **Busel et al.** <sup>(17)</sup>.

We did not make an attempt to recommend for or against fibular fixation as this was not our aim, controversy remains regarding when and how to perform the fibular fixation that is associated with pilon fractures in 80% of cases **Tomás-Hernández** <sup>(16)</sup>.

## Conclusion

The present study provides information on the use and the role of ankle arthroscopy in assisting and evaluation of distal tibial intra articular reduction after pilon fracture, also the value, advantages, techniques of ankle arthroscopy and ring external fixators. The study showed that this technique might be a good method to manage the pilon fractures.

The results reported in this study could contribute towards the importance of this combined technique and the encouraging outcome.

## References

1. **Bocchi L, Maniscalco P, Bertone C et al. (2000):** Fractures of the tibial plafond: a comparison of treatment methods. *Journal of Orthopaedics and Traumatology*, 1(1):51-56.
2. **Rüedi T, Allgöwer M (1969):** Fractures of the lower end of the tibia into the ankle-joint. *Injury*, 1:92-99.
3. **Ovadia D, Beals R (1986):** Fractures of the tibial plafond. *J Bone Joint Surg Am.*, 68:543-551.
4. **Association Committee for Coding and Classification. (1996):** Fracture and dislocation compendium. *J Orthop Trauma*, 10 (1):1-154.
5. **Blauth M, Bastian L, Krettek C (2001):** Surgical Options for the Treatment of Severe Tibial Pilon Fractures: A Study of Three Techniques. *Journal of Orthopaedic Trauma*, 15(3):153-60.
6. **Aggarwal A, Nagi O (2006):** Hybrid external fixation in periarticular tibial fractures: Good final outcome in 56 patients. *Acta Orthopaedica Belgica*, 72(4):434-40.
7. **Vidyadhara S, Rao S (2006):** Ilizarov treatment of complex tibial pilon fractures. *Int Orthop.*, 30(2): 113-117.
8. **Salter R, Simmonds D, Malcom B (1980):** The biological effect of continuous passive motion on the healing of full thickness defect in articular cartilage. An experimental investigations in the rabbit. *J Bone and Joint Surg Am.*, 26:1232-1251.
9. **El-Mowafi H, El-Hawary A, Kandil Y et al. (2015):** The management of tibial pilon fractures with the Ilizarov fixator: The role of ankle arthroscopy. *Foot (Edinb).*, 25:238-243.
10. **Lonjon G, Delgrande D, Solignac N et al. (2015):** Arthroscopic treatment in split depression-type tibial pilon fracture. *Eur J Orthop Surg Traumatol.*, 25(2):399-403.
11. **Zhao L, Li Y, Chen A et al. (2013):** Treatment of type C pilon fractures by external fixator combined with limited open reduction and absorbable internal fixation. *Foot Ankle Int.*, 34(4):534-42.
12. **Sirkin M, Sanders R, DiPasquale T et al. (2004):** A staged protocol for soft tissue management in the treatment of complex pilon fractures. *J Orthop Trauma.*, 18(8):32-8.
13. **Kapoor S, Kataria H, Patra S et al. (2010):** Capsuloligamentotaxis and definitive fixation by an ankle-spanning Ilizarov fixator in high-energy pilon fractures. *J Bone Joint Surg Br.*, 92(8):1100-6.

14. **Guillo S, Cordier G, Sonnery-Cottet B *et al.* (2014):** Anatomical reconstruction of the anterior talofibular and calcaneofibular ligaments with an all-arthroscopic surgical technique. *Orthop Traumatol Surg Res.*, 100(8):S413-7.
15. **Chiang C, Tzeng Y, Jeff Lin C *et al.* (2019).** Arthroscopic Reduction and Minimally Invasive Surgery in Supination-External Rotation Ankle Fractures: A Comparative Study With Open Reduction. *Arthroscopy*, 35(9):2671-2683.
16. **Tomás-Hernández J. (2017):** High-energy pilon fractures management: State of the art. *EFORT Open Rev.*, 1(10):354-361.
17. **Busel G, Watson J, Israel H (2017):** Evaluation of Fibular Fracture Type vs Location of Tibial Fixation of Pilon Fractures. *Foot Ankle Int.*, 38(6):650-655.