

Clinical and radiological manifestations of Central Nervous System (CNS) tuberculosis

¹Dr. MDinesh, ²Dr. Anith Alfred J, ³Dr. GSrirama Murthy,

¹Associate Professor, Department of Radiodiagnosis Shri Sathya Sai Medical College and Research Institute, Ammapettai, Chengalpattu, Tamil Nadu, India

²Assistant Professor, Department of Radiodiagnosis Shri Sathya Sai Medical College and Research Institute, Ammapettai, Chengalpattu, Tamil Nadu, India

³Professor, Department of Radiodiagnosis Shri Sathya Sai Medical College and Research Institute, Ammapettai, Chengalpattu, Tamil Nadu, India

Corresponding Author:

Dr. GSrirama Murthy (gurram1993@gmail.com)

Abstract

Background: Tuberculosis has resurged and remained a major worldwide health problem. Although Mycobacterium tuberculosis can involve any organ, most commonly the lung, central nervous system (CNS) tuberculosis is the most devastating form of the disease. Approximately 5-10% of all patients with tuberculosis and up to 20% of patients with AIDS-related tuberculosis have CNS involvement. Central nervous system (CNS) tuberculosis is a potentially life-threatening condition which is curable if the correct diagnosis is made in the early stages. Its clinical and radiologic manifestations may mimic other infectious and non-infectious neurological conditions. Hence, familiarity with the imaging presentations of various forms of CNS tuberculosis is essential in timely diagnosis, and thereby reducing the morbidity and mortality of this disease.

Aims and Objectives: To Evaluate the clinical and radiological manifestations of Central Nervous System (CNS) tuberculosis.

Material and Methods: Our study was conducted in Department of Radiodiagnosis Shri Sathya Sai Medical College and Research Institute, Ammapettai, Chengalpattu, Tamil Nadu, India and to describe and get familiarized with the imaging characteristics of the different forms of CNS tuberculosis in 50 indoor patients of tuberculosis diagnosed by techniques involving culture and immunological testing of the tissue and biofluids and referred to the department of radio-diagnosis for imaging. Patients with spinal tuberculosis were excluded in our study.

Results and Observations: Males and females contributed 46% and 54% of the study population respectively and in. The age range in years was 10 to 76 years. The mean age of the patients was 28 years with 60% of the patients between 20-40 years. Out of the 50 patients, the commonest presentation was meningitis (82%) especially seen in the younger population-children and adolescents. Early diagnosis of CNS TB is necessary for appropriate treatment and subsequently decreases the morbidity and mortality associated with the disease. Routine diagnostic techniques involve culture and immunological tests of the tissue and biofluids, which are time-consuming and may delay definitive management.

Conclusion: CNS tuberculosis has various imaging appearances, including meningitis, tuberculoma, miliary tuberculosis, abscess, cerebritis, and encephalopathy. In addition, the radiologic manifestations of this disease are not always typical and sometimes may be mistaken with other lesions. Familiarity with the various imaging presentations of CNS tuberculosis is of key importance for the radiologists and infectious diseases specialists in timely diagnosis, thereby reducing the morbidity and mortality of this potentially life-threatening disease. Non-invasive imaging modalities such as computed tomography (CT) scan and magnetic resonance imaging (MRI) are routinely used in the diagnosis of neurotuberculosis, with MRI offering greater inherent sensitivity and specificity than CT scan. In this review, we describe the imaging characteristics of the different forms of CNS tuberculosis, including meningitis, tuberculoma, miliary tuberculosis, abscess, cerebritis, and encephalopathy.

Keywords: CT Scan, magnetic resonance imaging, CNS tuberculosis, radiological manifestations, non-invasive imaging, early diagnosis, miliary tuberculosis

Introduction

Infection of CNS by tuberculosis (TB) can develop in several patterns. The old adage “Syphilis-the Great Masquerader”, probably applies to tuberculosis as well when it involves the central nervous system. This poses a constant challenge to the clinicians in making a definitive diagnosis. The various extra-pulmonary manifestations of tuberculosis and the multitude of organs involved complicate the task further. Central nervous system (CNS) tuberculosis forms a significant proportion of extra-pulmonary tuberculosis. Mycobacteria are small rod-shaped acid-fast bacilli with more than 125 species. Most TB is caused by mycobacterium tuberculosis. It causes more than 98% of the CNS tuberculosis. Less common species that are also considered part of the M. tuberculosis complex include M. africanum, M. microti, M. canetti and M. bovis. Tuberculosis has resurged and remained a major worldwide health problem. Although Mycobacterium tuberculosis can involve any organ, most commonly the lung, central nervous system (CNS) tuberculosis is the most devastating form of the disease. Approximately 5-10% of all patients with tuberculosis and up to 20% of patients with AIDS-related tuberculosis have CNS involvement^[1, 3]. CNS tuberculosis usually results from hematogenous spread, while direct spread from intra-or extracranial focus is rare^[4]. The clinical and radiologic manifestations of CNS tuberculosis may mimic other infectious and non-infectious neurological conditions. Therefore, familiarity of infectious diseases specialists with the imaging presentations of CNS tuberculosis is essential for prompt and accurate diagnosis of this entity. Herein, we describe the different forms of CNS tuberculosis including meningitis, cerebritis, cerebral abscesses, tuberculomas, miliary tuberculosis and spinal or calvarial involvement^[5, 7].

Material and Methods

Our study was conducted in Department of Radiodiagnosis Shri Sathya Sai Medical College and Research Institute, Ammapettai, Chengalpet, TamilNadu, India and to describe and get familiarized with the imaging characteristics of the different forms of CNS tuberculosis in 50 indoor patients of tuberculosis diagnosed by techniques involving culture and immunological testing of the tissue and biofluids and referred to the department of radio-diagnosis for imaging. Patients with spinal tuberculosis were excluded in our study.

Results and Observations


There are four major patterns of CNS TB:

1. Tuberculous meningitis (TBM).
2. Tuberculomas in brain and spinal cord (TBT).
3. Tubercular brain abscess (TBA).

4. Tuberculous encephalopathy (TBE).

Table 1: Sex distribution

Sex	No of patients	Percentage(%)
Males	23	46
Females	27	54
Total	50	100

**Fig 1:** Sex distribution

Males and females contributed 46% and 54% of the study population respectively and in Table 1 and Figure 1. The age range in years was 10 to 76 years. The mean age of the patients was 28 years with 60% of the patients between 20-40 years. Out of the 50 patients, the commonest presentation was meningitis (82%) especially seen in the younger population - children and adolescents.

Discussion

Surgery has a role both in the diagnosis and treatment of tuberculoma. It is usually indicated in cases of clinical deterioration that fail to respond to medical management and when the diagnosis of tuberculoma is in doubt. Surgical treatment options for TBA include simple puncture, continuous drainage, fractional drainage, repeated aspiration through a burr hole, stereotactic aspiration and total excision of the abscess. Ventriculoperitoneal shunts may be required in patients with TBM or TBT who develop obstructive hydrocephalus. Ventriculoperitoneal shunt should be performed at the time the need for shunting is determined and should not be delayed waiting for the infection to resolve.

Tuberculous meningitis (TBM) is the most common cause of chronic meningitis, especially in developing countries. Meningitis is the most common manifestation of CNS tuberculosis which is most frequently seen in the children and adolescents^[8, 9]. Tuberculous meningitis is mostly due to the hematogenous spread of *Mycobacterium tuberculosis*^[10]. Tuberculous meningitis often has an insidious course with a nonspecific clinical presentation in early stages, especially in children. Therefore, the imaging plays a key role in the timely diagnosis and decreasing the morbidity and mortality. TB meningitis has a striking predilection for the

basal cisterns although exudates in the superficial convexity sulci do occur. Enhancing exudate in the basal cisterns is the most common and also a relatively specific manifestation of leptomeningeal tuberculosis on computed tomography (CT) and magnetic resonance (MR) images. Meningeal enhancement has been found in up to 90% of cases and is considered to be the most sensitive feature of tubercular meningitis^[11, 12]. The subpial exudate is primarily located in the inferomedial surface of the frontal lobes, the anteromedial surface of the temporal lobes, the superior aspect of the cerebellum, and the floor of the third ventricle^[13]. Extension to suprasellar, interpeduncular, and pontomesencephalic cisterns may also occur

from these primary sites. In most cases, some degree of meningeal involvement is seen within the sulci over the cerebral convexities, the sylvian fissures and also the ependymal surfaces of the ventricles; the latter usually occurs in the later stages of the disease^[14, 15]. CT findings; CT can be normal in early stages. On CT images, the obliteration of the basal cisterns by isodense or mildly hyperdense exudates is the most common finding in tuberculous meningitis^[16]. “blurred” ventricular margins may be seen.

As in Figure 2-6, The findings are better appreciated on MR imaging than on CT, especially on postcontrast MR images which show the enhancing cisternal exudates and leptomeningeal enhancement. Basilar exudates are isointense with brain on T1WI giving the appearance of “dirty” CSF. FLAIR scans show increased signal intensity in the sulci and cisterns. Marked linear or nodular meningeal enhancement is seen on T1 C + FS sequences. Focal or diffuse dura-arachnoid enhancement (pachymeningeal) with or without involvement of the underlying subarachnoid spaces may occur but is uncommon. Parmar *et al.* demonstrated that postcontrast fluid attenuation inversion recovery (FLAIR) images may have a higher specificity compared to contrast-enhanced T1-weighted images in detection of leptomeningeal enhancement^[17]. Magnetization transfer spin echo imaging following contrast injection is superior to the conventional postcontrast imaging in demonstrating meningeal inflammation. In later stages, there may be widening of subarachnoid spaces. A similar pattern of meningeal enhancement may be seen in other infective meningitis, inflammatory diseases such as rheumatoid arthritis, sarcoidosis, or carcinomatous meningitis. Carcinomatous meningitis is usually seen in older patients with a known systemic or primary CNS neoplasm. In neurosarcoidosis, infiltration of the pituitary gland, infundibulum and hypothalamus is common. Other radiologic manifestations of tuberculous meningitis may be related to its possible complications, including progressive hydrocephalus, vasculitis, infarction, and cranial neuropathies.

Hydrocephalus is the most common complication encountered in TBM and can be broadly divided into two types: communicating type, which is common, secondary to an obstruction of the basal cisterns by inflammatory exudates and obstructive type, which is less common and either secondary to a focal parenchymal lesion (tuberculoma or a brain abscess) causing mass effect or due to the entrapment of a part of the ventricle by granulomatous ependymitis. It is a complication that is commonly seen at autopsy in cranial TBM. The adventitial layer of small and medium-sized vessels develops changes similar to those of the adjacent tuberculous exudates. The intima of the vessels may eventually be affected or eroded by fibrinoid-hyaline degeneration. In later stages, the lumen of the vessel may get completely occluded by reactive subendothelial cellular proliferation. Ischemic cerebral infarction resulting from the vascular occlusion is a common sequelae of tuberculous arteritis. Ischemic infarct is also a common complication, being detected in 20-41% of patients on CT, mostly within the basal ganglia or internal capsule regions and resulting from vascular compression and occlusion of small perforating vessels (necrotizing arteritis)^[18, 19], particularly the lenticulostriate and thalamoperforating arteries, vessels which perfuse the so-called medial tuberculosis zone¹⁶. Tuberculous meningitis may also cause dural venous sinus thrombosis with resultant hemorrhagic infarct. Rarely, tuberculosis may present as isolated dural venous sinus thrombosis without any evidence of meningitis or its complications. Cranial nerve involvement is associated with high morbidity and disability in meningitis patients. Cranial nerve involvement occurs due to vascular compromise, ischemia, or nerve entrapment in the basal exudates in 17-40% of cases, most commonly affecting the second, third, fourth, and seventh cranial nerves. The affected cranial nerves are best evaluated by MRI, where they may appear thickened, especially in their proximal segments, with high signal intensity on T2-weighted images and marked enhancement on postcontrast images.

Parenchymal Tuberculosis: Parenchymal disease may be isolated or associated with tuberculous meningitis. Parenchymal involvement usually presents as tuberculoma. It can also manifest as cerebritis, cerebral abscess, miliary tuberculosis, or tuberculosis encephalopathy^[20, 21, 22].

A. Cerebritis and Cerebral Abscess

Parenchymal tuberculosis may occur with or without accompanying meningitis. Tuberculosis cerebritis or abscess may have an appearance similar to that of pyogenic bacterial infection on neuroimaging studies. Focal tuberculous cerebritis is very rare, cerebritis is a term that represents inflammation of the brain in the setting of infection before the development of abscess. CT: Hypoattenuation due to edema is seen. Small areas of hemorrhage can sometimes be seen. Faint heterogeneous contrast enhancement can be seen. MRI: T1 iso to hypointensity, T2 hyperintensity, and minimal post contrast enhancement on T1 C + images is seen. Patchy restricted diffusion is seen. Tuberculous pseudo-abscess is rare and is characterized by a central area of liquefaction with pus. It may be solitary or multiple and is frequently multiloculated. Tuberculous abscess is different from tuberculomas which contain central caseation and liquefaction mimicking pus. On CT: Tb pseudoabscess are hypodense on NECT with significant mass effect and surrounding edema. Ring enhancement is seen on CECT. On MRI: Unlike tuberculomas, Tb pseudoabscess are usually hyperintense to brain on T2/FLAIR images and restrict on DWI. A ring enhancing multi-loculated lesion or multiple separate lesions is the typical finding on T1 C + images. MRS shows lipid and lactate peaks without evidence of cytosolic amino-acids. TB pseudoabscess appear identical to pyogenic abscesses on standard imaging studies. Both show restricted diffusion. On MR spectroscopy, the peak of amino acids, which is the spectral hallmark in pyogenic abscess, is not usually seen in tuberculous pseudoabscess^[23-27]. Magnetization transfer (MT) images improve the conspicuity of all CNS tuberculosis lesions.

B. Tuberculoma

Tuberculoma is the most common parenchymal lesion in CNS tuberculosis which could be found in any portion of the intracranial space. The lesion may be solitary or multiple and may be seen with or without meningitis. Histologically, the mature tuberculoma is composed of a necrotic caseous center surrounded by a capsule that contains fibroblasts, epithelioid cells, Langhans giant cells and lymphocytes^[28]. On CT: NECT scans show one or more iso to slightly hyperdense round, lobulated or crenated masses with variable perilesional edema. Calcification can be seen in healed granulomas. CECT scans demonstrate punctate, solid or ring like enhancement. MRI: Most TB granulomas are solid caseating lesions that appear hypo or isointense with brain on T1WI and hypointense on T2WI. Liquefied areas may be T2 hyperintense with a hypointense rim. Enhancement is variable ranging from small punctate foci to multiple rim enhancing lesions. Mild to moderate round or lobulated ring like enhancement around a non-enhancing center is the most typical pattern. pMR shows elevated rCBV in the cellular, hypervascular enhancing rim. Solid caseating tuberculomas do not restrict on DWI although liquefied foci may restrict. MRS is helpful in characterizing

tuberculomas and distinguishing them from neoplasm or pyogenic abscess. A prominent decrease in the NAA: Cr with a modest decrease in the NAA: Cho is typical. A large lipid peak with absence of other metabolites such as amino acids and succinate is seen in 85-90 % of the cases. The major differential for multiple parenchymal tuberculomas is seen in neurocysticercosis (NCC). NCC usually shows multiple lesions in different stages of evolution. Tuberculomas can also resemble pyogenic abscesses or neoplasms. Abscesses restrict on DWI. Tuberculomas have a large lipid peak on MRS and lack the elevated Cho typical of neoplasm^[28]. Follow-up CT or MR studies are useful in monitoring the response to medical treatment. Paradoxical enlargement of a pre-existing tuberculoma or evolution of a new intracranial and spinal tuberculoma in patients receiving adequate treatment may be occasionally seen. However, with continuation of anti-tuberculous therapy, eventual resolution of the tuberculoma usually occurs^[29,30]. Sometimes, healed tuberculomas appear as calcified foci on nonenhanced CT. Similarly, calcification in the basal cisterns has been demonstrated a

few years after meningeal tuberculosis^[31].

C. Miliary tuberculosis

Miliary tuberculosis is seen mostly in severely immune compromised patients and is usually associated with meningeal involvement or extracranial primary sites^[32]. Since the dissemination is hematogenous, the lesions are usually located at the corticomedullary junctions. The lesions are tiny (2-3 mm in diameter) scattered lesions that may be invisible on non-contrast MR sequences in visible lesions, MRI shows small lesions that are hypointense on T2-weighted sequences. These lesions occasionally can be seen as small hypodensities on CT scan. Postcontrast T1-weighted MR images show numerous, round, small, homogeneous, enhancing (usually ring enhancement) lesions²². Invisible lesions that may or may not enhance after intravenous injection of gadolinium can be clearly visible on magnetization transfer spin echo T1-weighted imaging with or without contrast.

D. Tuberculous encephalopathy

Tuberculous encephalopathy typically occurs in young children who may present with convulsion, stupor and coma with no signs of meningeal irritation or focal neurological deficit. Neuroimaging studies show severe cerebral edema, which may be unilateral or bilateral. Myelin loss in the white matter may result in hypodensity on CT images and hyperintensity on T2-weighted MR images.

Osseous and non-osseous spinal/spinal cord tuberculosis, subdural/epidural abscess and calvarial tuberculosis are other forms of tuberculosis that may involve CNS with direct or indirect pathways. Tuberculous spinal meningitis presents on MR imaging as a CSF loculation and obliteration of the spinal subarachnoid space, with loss of the outline of the spinal cord in the cervicothoracic spine and matting of the nerve roots in the lumbar region. Contrast-enhanced imaging reveals nodular, thick, linear intradural enhancement, which may completely fill the subarachnoid spaces. Longstanding arachnoiditis may result in the development of syringomyelia (spinal cord cavitation) that typically demonstrates CSF signal intensity on all MR sequences. Tuberculous spondylitis results from hematogenous spread of infection to the vertebral body via paravertebral venous plexus of Batson. Typical presentation is involvement of multiple vertebral bodies with sparing of intervertebral disc in early stage and disc involvement in later stages. Paraspinal extension and resultant paravertebral abscess (Pott's abscess) as well as subdural/epidural abscess formation with associated spinal cord compression are other common findings. Intracranial subdural or epidural abscess formation may or may not be associated with a primary CNS tuberculous focus and have imaging findings identical to that of other pyogenic abscesses, that is, iso-to hypointensity on T1-weighted images, hyper- or mixed signal intensity on T2-weighted images, and rim enhancement on postcontrast images.




Fig2: T2W axial, T2W FLAIR axial, T1W C +,SWI, ADC and DWI images showing tuberculomas in the right frontal lobe




Fig3: T1W axial, T2W axial, T1W C +, DWI, ADC and T2 C+ FLAIR images showing right cerebellar tuberculoma. Vasculitic infarct in right thalamic-capsular region. Extensive basilar meningitis. Post-operative defect with meningoencephalocele is also noted involving the right occipital lobe and right cerebellar hemisphere at the post-operative site




Fig4: T2W FLAIR axial images in the same patient showing hydrocephalus




Fig5: CT brain plain and delayed post contrast axial images showing a small sub centimeter sized ring enhancing lesion in the right parietal sub-cortical white matter with mild adjacent edema in this patient with past history of sputum positive pulmonary tuberculosis. The patient presented with convulsions



Fig6: CT brain plain axial image showing a calcified granuloma in the genu of corpus callosum in a case of RVD patient with PMLE and TB

Conclusion

Central nervous system tuberculosis is one of the more severe forms of extra-pulmonary tuberculosis. There has been a significant increase in the incidence of CNS TB worldwide

over the last couple of decades. It may present as tuberculous meningitis, tuberculoma, tuberculous abscess, or encephalopathy. Its diagnosis relies on. CNS tuberculosis has various imaging appearances, including meningitis, tuberculoma, miliary tuberculosis, abscess, cerebritis, and encephalopathy. In addition, the radiologic manifestations of this disease are not always typical and sometimes may be mistaken with other lesions. Familiarity with the various imaging presentations of CNS tuberculosis is of key importance for the radiologists and infectious diseases specialists in timely diagnosis, thereby reducing the morbidity and mortality of this potentially life-threatening disease.

laboratory studies of the CSF and/or visualization of a lesion on CT or MRI. Newer diagnostic techniques, such as PCR, have not been assessed completely and are not possible in most settings in the developing world. Neurosurgeons have been actively involved in the treatment of central nervous system tuberculosis due to its propensity to cause obstructive hydrocephalus, intracranial mass lesions, and compressive myelopathy. Surgery has a role both in the diagnosis and/or treatment of tuberculoma and tubercular brain abscess although definite guidelines have yet to be formulated. A high index of suspicion is required in order to avoid delays in diagnosis, which may influence treatment outcome. Death may occur as a result of missed diagnosis and delayed treatment.

Source of Support: None.

Conflict of Interest: None.

References

1. Bernaerts A, Vanhoenacker FM, Parizel PM, *et al.* Tuberculosis of the central nervous system: overview of neuroradiological findings. *European Radiology*. 2003;13(8):1876-1890.
2. Berenguer J, Moreno S, Laguna F, *et al.* Tuberculous meningitis in patients infected with the human immunodeficiency virus. *The New England Journal of Medicine*. 1992;326(10):668-672.
3. Vidal JE, De Oliveira ACP, Filho FB, *et al.* Tuberculous brain abscess in AIDS patients: report of three cases and literature review. *International Journal of Infectious Diseases*. 2005;9(4):201-207.
4. Rock RB, Olin M, Baker CA, Molitor TW, Peterson PK. Central nervous system tuberculosis: pathogenesis and clinical aspects. *Clinical Microbiology Reviews*.

2008;21(2):243-261.

5. Burrill J, Williams CJ, Bain G, Conder G, Hine AL, Misra RR. Tuberculosis: a radiologic review. *Radiographics*. 2007;27(5):1255-1273.
6. Bathla G, Khandelwal G, Maller VG, Gupta A. Manifestations of cerebral tuberculosis. *Singapore Medical Journal*. 2011;52(2):124-131.
7. Ahluwalia VV, Dayananda Sagar G, Singh TP, Arora N, Narayan S, Singh MM. MRI spectrum of CNS tuberculosis. *Journal, Indian Academy of Clinical Medicine*. 2013;14(2):83-90.
8. Raviglione MC, Snider DE Jr, Kochi A. Global epidemiology of tuberculosis: morbidity and mortality of a worldwide epidemic. *The Journal of the American Medical Association*. 1995;273(3):220-226.
9. Andronikou S, Wilmshurst J, Hatherill M, VanToorn R. Distribution of brain infarction in children with tuberculous meningitis and correlation with outcome score at 6 months. *Pediatric Radiology*. 2006;36(12):1289-1294.
10. Dastur DK, Manghani DK, Udani PM. Pathology and pathogenetic mechanisms in neurotuberculosis. *Radiologic Clinics of North America*. 1995;33(4):733-752.

11. Andronikou S,Smith B,Hatherhill M,Douis H,Wilmshurst J. Definitive neuroradiological diagnostic features of tuberculous meningitis in children. *Pediatric Radiology*. 2004;34(11):876-885.
12. Uysal G,Köse G,Güven A,Diren B. Magnetic resonance imaging in diagnosis of childhood central nervous system tuberculosis. *Infection*. 2001;29(3):148-153.
13. Arbeláez A,Medina E,Restrepo F,Castillo M. Cerebral tuberculosis. *Seminars in Roentgenology*. 2004;39(4):474-481.
14. Jinkins JR,Gupta R,Rodriguez-Carbajal J. MR imaging of central nervous system tuberculosis. *Radiologic Clinics of North America*. 1995;33(4):771-786.
15. Morgado C,Ruivo N. Imaging meningo-encephalic tuberculosis. *European Journal of Radiology*. 2005;55(2):188-192.
16. Shah GV. Central nervous system tuberculosis: imaging manifestations. *Neuroimaging Clinics of North America*. 2000;10(2):355-374.
17. Parmar H,Sitoh YY,Anand P,Chua V,Hui F. Contrast-enhanced flair imaging in the evaluation of infectious leptomenigeal diseases. *European Journal of Radiology*. 2006;58(1):89-95.
18. Dastur DK,Lalitha VS,Udani PM,Parekh U. The brain and meninges in tuberculous meningitis-gross pathology in 100 cases and pathogenesis. *Neurology India*. 1970;18(2):86-100.
19. Whiteman MLH. Neuroimaging of central nervous system tuberculosis in HIV-infected patients. *Neuroimaging Clinics of North America*. 1997;7(2):199-214.
20. Sharma P, Garg RK,Verma R,Singh MK,Shukla R. Incidence, predictors and prognostic value of cranial nerve involvement in patients with tuberculous meningitis: a retrospective evaluation. *European Journal of Internal Medicine*. 2011;22(3):289-295. Doi: 10.1016/j.ejim.2011.01.007.
21. Garg RK. Tuberculosis of the central nervous system. *Postgraduate Medical Journal*. 1999;75(881):133-140.
22. Trivedi R,Saksena S,Gupta RK. Magnetic resonance imaging in central nervous system tuberculosis. *Indian Journal of Radiology and Imaging*. 2009;19(4):256-265. Doi: 10.4103/0971-3026.57205.
23. Gupta RK,Husain N,Kathuria MK,Datta S,Rathore RKS,Husain M. Magnetization transfer MR imaging correlation with histopathology in intracranial tuberculomas. *Clinical Radiology*. 2001;56(8):656-663.
24. Saxena S, Prakash M, Kumar S, Gupta RK. Comparative evaluation of magnetization transfer contrast and fluid attenuated inversion recovery sequences in brain tuberculoma. *Clinical Radiology*. 2005;60(7):787-793.
25. Gupta RK, Kathuria MK,Pradhan S. Magnetization transfer MR imaging in CNS tuberculosis. *The American Journal of Neuroradiology*. 1999;20(5):867-875.
26. Gupta R. Magnetization transfer MR imaging in central nervous system infections. *Indian Journal of Radiology and Imaging*. 2002;12(1):51-58.
27. Gupta RK,Vatsal DK,Husain N, *et al*. Differentiation of tuberculous from pyogenic brain abscesses with *in vivo* proton MR spectroscopy and magnetization transfer MR imaging. *American Journal of Neuroradiology*. 2001;22(8):1503-1509.
28. Kim TK,Chang KH,Kim CJ,Goo JM,Kook MC,Han MH. Intracranial tuberculoma: comparison of MR with pathologic findings. *The American Journal of Neuroradiology*.1995;16(9):1903-1908.
29. Ku BD, Yoo S. Extensive meningeal and parenchymal calcified tuberculoma as long-term residual sequelae of tuberculous meningitis. *Neurology India*. 2009;57(4):521-522.
30. Afghani B,Lieberman JM. Paradoxical enlargement or development of intracranial tuberculomas during therapy: case report and review. *Clinical Infectious Diseases*. 1994;19(6):1092-1099.

31. Lorber J. Intracranial calcifications following tuberculous meningitis in. *Acta Radiologica*. 1958;50(1-2):204-210.
32. Krishnan N, Robertson BD, Thwaites G. The mechanisms and consequences of the extra-pulmonary dissemination of *Mycobacterium tuberculosis*. *Tuberculosis*. 2010;90(6):361-366.
33. Udani PM, Dastur DK. Tuberculous encephalopathy with and without meningitis clinical features and pathological correlations. *Journal of the Neurological Sciences*. 1970;10(6):541-561.