

Original research article

Assessment Of The Expenses And Benefits Of Procalcitonin Testing In The Diagnosis Of Early Onset Neonatal Sepsis**Dr. Manjul Vijay¹, Dr. Tejaswi Nandan²****¹Senior Resident, Department of Paediatrics, Anugrah Narayan Magadh Medical College and Hospital, Gaya, Bihar, India****²Senior Resident, Department of Obstetrics and Gynecology, Anugrah Narayan Magadh Medical College and Hospital, Gaya, Bihar, India.****Corresponding Author: Dr. Tejaswi Nandan****Abstract**

Aim: To compare between the expenses and benefits of procalcitonin testing in the diagnosis of early onset neonatal sepsis.

Material and methods: This prospective observational study was carried out in the Department of Pediatrics, Anugrah Narayan Magadh medical college and hospital Gaya, Bihar, India for one year. 100 Full-term neonates (>37 weeks gestational age) admitted to NICU with the clinical symptoms or signs of sepsis within 72 h of birth, and those with risk factors for EONS. For all patients included in the study, complete blood count with differential (CBC), CRP, blood culture, procalcitonin was done on admission to NICU, and was repeated after 8 hours from the initial one, while CRP was repeated after 24 hrs. Cultures from other sites (including CSF), chest x ray and imaging were done as appropriate.

Results: 100 patients were included in the study, 70 females and 30 males. Their mean weight was 3.163 kg (range 2.1- 4.9 kg). About 89 (89 %) patients presented with more than one symptoms and signs of sepsis. For CBC results, mean WBC count was 19.9 ± 10.1 , the mean neutrophils count was 15.45 ± 11.57 and the mean platelet count was 168.21 ± 70.21 . The most frequent presenting sign was hypothermia (62%), hypoactivity and mottling (53%), followed by feeding intolerance (42%). Blood cultures were positive in 35 patients, who compromised the group of proven sepsis. Klebsiella was present in 13 patient, E coli in 9 patients, while group B streptococcus in 8 patients. Pseudomonas infection present in 3 patients, while MRSA (methicillin resistant staphylococcus aureus) was detected in 2 patients. The initial CRP (CRP1), was non-reactive in 45 patients (45%), The initial procalcitonin in the sepsis group (PCT1), was non-reactive 45 patients (45%), <2.6 in 31 patients (31%), 2.6-10 in 19 patients (19%), and more than that in 5 patients (5%). The 35 patients in the group of proven infection (35%) were admitted for 10-14 days but 5 of them needed more than 2 WEEKS admission. The second group (not-proven infection) of 45 patients (45%) were discharged within 4 days of admission, and they did well on the follow up except for 8 patients who didn't show up. The third group 20 patients (20%) suspected infection, treated with antibiotics for at least 10 days, and all of them did well on the follow up. CRP2 Sensitivity (92.5%), Specificity(53%), PPV(65.3%), NPV(86.5%) and PCT2 Sensitivity(95.8%), Specificity(87.7%), PPV(90.1%), NPV(98%).

Conclusion: The expense of testing PCT twice is less than the expenses of one-day admission in NICU in developing countries. Application of such protocol could be of use in limiting period of stay in NICU.

Introduction

Neonatal sepsis (NS), a life-threatening emergency, a 3rd major cause for mortality¹ and morbidity globally, remains a concern despite the breakthroughs in the diagnosis and

management. In India, the estimated incidence of NS is 30/1000 live births contributing to 19% of all neonatal deaths.² The symptoms and signs of NS mimicking that of other infections, co-existence of infections make the diagnosis difficult, particularly in countries battling with infectious diseases. With varying incidence in Asian countries, and \approx 1.6 million neonates succumbing annually in developing countries indicates the need for an early diagnosis including clinical and laboratory evaluation.³ There are many available markers for NS but each with limitations; Complete Blood Count (CBC), immature: total neutrophil count (I: T ratio) and absolute neutrophil count (ANC) do not have sensitivity especially if measured early in the course of sepsis.⁴ Though culture is the gold standard for confirmation,⁵ result is influenced by various factors such as prior use of antibiotics including in the prenatal period, bacterial load, laboratory standards and is time-consuming (up to 72 hours); it often fails to identify the causative organism in infected infants⁶ given low culture yields. Several leukocyte indices and acute phase protein levels were evaluated for the diagnosis of sepsis. C-reactive protein though is a classical and sensitive marker of inflammation⁷ is not useful to differentiate between bacterial and other infections, has a limited role in diagnosing early-onset sepsis compelling the use of combination tests for markers.⁸ In the absence of a single ideal diagnostic & confirmatory laboratory test,⁷ the quest and search for the same for early diagnosis of NS is the need of the hour. The paradigm in pediatric practice is a neonate is likely to suffer more if the infection is under-diagnosed and untreated than over-diagnosed and treated demanding a diagnostic test with high sensitivity than high specificity. Emerging shreds of evidence showing serum procalcitonin (PCT) as a measurable laboratory marker in the inflammatory response to the infection is promising due to its high sensitivity compared to CRP, but its specificity is still debated.⁹ Considering the diversity in population, that is different from other global counterparts where the diagnostic utility of PCT is encouraging, there is a scarcity of studies on evaluation of the role of various biochemical markers in the diagnosis of NS especially PCT in Indian neonates. We attempted to document the effects of intrapartum risk factors, assess and compare the diagnostic role of Procalcitonin and CRP in early onset NS.

Material and methods

This prospective observational study was carried out in the Department of Pediatrics, Anugrah Narayan Magadh medical college and hospital Gaya, Bihar, India for one year. after taking the approval of the protocol review committee and institutional ethics committee.

Inclusion criteria

100 Full-term neonates (>37 weeks gestational age) admitted to NICU with the clinical symptoms or signs of sepsis within 72 h of birth, and those with risk factors for EONS. The risk factors considered were: GBS infection during pregnancy, premature rupture of membrane, prolonged rupture >18 h before birth and Clinical syndrome of maternal intrauterine infection.

Exclusion criteria

Patients < 37 weeks gestational age, patients with congenital anomaly or metabolic inborn error of metabolism, patients received antibiotics before admission to NICU and refusal of the parents to sign the consent.

Methodology

For all patients included in the study, complete blood count with differential (CBC), CRP, blood culture, procalcitonin was done on admission to NICU, and was repeated after 8 hours

from the initial one, while CRP was repeated after 24 hrs. Cultures from other sites (including CSF), chest x ray and imaging were done as appropriate.

Under complete aseptic conditions 3 ml of venous blood was collected, 1 ml on ethylene diamine tetra acetic acid (EDTA) for complete blood count (CBC) and 2 ml was collected in plane tube and was left to clot then centrifuged to separate serum for estimation of serum CRP and detection of PCT. For serum PCT analysis Sera of the patients were analyzed for PCT using commercially available Enzyme-linked immunosorbent assay (ELISA) kit. PCT production was calculated from a standard curve of the corresponding recombinant human PCT. Manual broth-based blood culture systems was used, namely non-selective agar media. Growth of any organism in samples taken from symptomatic newborns was taken significant.

Criteria for the diagnosis of neonatal infection¹⁰

Temperature instability, Heart rate >180 beats/min or <100 beats/min, blood pressure 2 SD below normal for age, and capillary refill >3s. Respiratory problems: as apnea, dyspnea, retractions and cyanosis. Central nervous system affection: irritability, lethargy, abnormal Moro reflex, fontanel bulging, seizures, and hypotonia. Gastrointestinal system affection as feeding intolerance, abdominal dis- tension with repeated vomiting or frequent watery motions. leukocytosis (WBC>34000), leukopenia (WBC<5000), Thrombocytopenia<100000, CRP 2 SD above normal level, Procalcitonin 2 SD above normal value By the end of the first 48 h newborns were labeled proven infected if their blood culture showed growth of organism; and were continued on antibiotics. Another group were labeled infected but not proven and those include patients with positive CRP, leukopenia or leukocytosis, or if have clinical symptoms and signs of sepsis. The third group is the rest of the newborn suspected for EONS but without evidence, they were labeled suspected only, and their antibiotics were discontinued.

Results

100 patients were included in the study, 70 females and 30 males. Their mean weight was 3.163 kg (range 2.1- 4.9 kg). Mean age of presentation was 14 hours (range 2-70) hours. 80 newborns (80%) presented within 28h of age, out of whom 52 newborns admitted at birth due to prolonged rupture of membrane (18 h and more). Prolonged rupture of membrane was the main risk factor, followed by maternal infections and fever (28 patients), fetal tachycardia (14 patients), and smelly liquor (6 patients).

Table 1 shows the clinical features of the newborns on admission to NICU About 89 (89 %) patients presented with more than one symptoms and signs of sepsis For CBC results, mean WBC count was 19.9 ± 10.1 , the mean neutrophils count was 15.45 ± 11.57 and the mean platelet count was 168.21 ± 70.21 . This study includes neonates with early onset sepsis. In the present study the most frequent presenting sign was hypothermia (62%), hypoactivity and mottling (53%), followed by feeding intolerance (42%).

Table 1: clinical symptoms and signs of the patients

Clinical symptoms and signs	Number of patients	Percentage
Tachypnea	36	36
Tachypnea with intercostal retractions	19	19
Sudden desaturation	13	13
Tachycardia	7	7
Bradycardia	16	16
Feeding intolerance, vomiting	42	42
	11	11

Hypothermia	62	62
Hypo activity, mottling	53	53

Blood cultures were positive in 35 patients, who compromised the group of proven sepsis. Klebsiela was present in 13 patient, E coli in 9 patients, while group B streptococcus in 8 patients. Pseudomonas infection present in 3 patients, while MRSA (methicillin resistant staphylococcus aureus) was detected in 2 patients. The initial CRP (CRP1) , was non-reactive in 45 patients (45%), The initial procalcitonin in the sepsis group (PCT1), was non-reactive 45 patients (45%) , <2.6 in 31 patients (31%), 2.6-10 in 19 patients (19%), and more than that in 5 patients (5%).

Table (2) shows CRP1/PCT1 results in the group with proven sepsis and those infected but not proven in Table (3). CRP2/PCT2 for the two categories are shown in Table 4 and 5. Table 6 shows work out of the predictive values of CRP2 and PCT2.

Table 2: Results of CRP1 and PCT1 in newborns with proven sepsis

	Positive	Negative	Total
CRP1	25	10	35
PCT1	27	8	35
Total	52	18	70 Pvalue 0.374

Table 3: Results of CRP1 and PCT1 in newborns with not-proven sepsis

	Positive	Negative	Total
CRP1	7	33	40
PCT1	4	36	40
Total	11	69	80 Pvalue 0.074

Table 4: Results of CRP2 and PCT2 in newborns with not-proven sepsis

	Positive	Negative	Total
CRP2	31	4	35
PCT2	32	3	35
Total	63	7	70 Pvalue 0.069

Table 5: Results of CRP2 and PCT2 in newborns with not-proven sepsis.

	Positive	Negative	Total
CRP2	20	20	40
PCT2	5	35	40
Total	25	55	80 Pvalue 0.003

Table 6: The predictive value for CRP2 and PCT2 in diagnosis of EONI.

	Positive	Treated group	Not treated group	Total
CRP2	Positive	55	23	78
	Negative	4	18	22
PCT2	Positive	51	4	55
	Negative	6	39	45

CRP2: Sensitivity= 92.5%, Specificity=53%, PPV= 65.3%, NPV= 86.5%. PCT2: Sensitivity=95.8%, Specificity=87.7%, PPV=90.1%, NPV=98%.

Managing infants in the three groups (proven, not-proven, and suspected infection) resulted in 411 inpatient days. The 35 patients in the group of proven infection (35%) were admitted for

10-14 days but 5 of them needed more than 2 weeks admission. four patients of this group died due to infection with GBS which caused bacteraemia, followed by toxic myocarditis and heart failure, that didn't respond to the antibiotic course. The second group (not-proven infection) of 45 patients (45%) were discharged within 4 days of admission, and they did well on the follow up except for 8 patients who didn't show up. The third group 20 patients (20%) WITH suspected infection, treated with antibiotics for at least 10 days, and all of them did well on the follow up.

Discussion

This study includes neonates with early onset sepsis. In the present study the most frequent presenting sign was hypothermia (62%), hypoactivity and mottling (53%), followed by feeding intolerance (42%). Mamta et al,¹¹ reported refusal to feed (77%), respiratory distress (44%), and hypothermia (47.5%), while Khatua et al.,¹² re-ported refusal to feed (92%), lethargy (74%), hypothermia (72%) and respiratory distress (24%) as common clinical presentation. In agreement with the study done by Muhammed et al, the most frequent recognized risk factor in the present study, was premature rupture of membrane (PROM).¹³

A study from Bangladesh showed that approximately one-third of all septicemia in neonates was attributable to premature rupture of membranes.¹⁴ While a study from Thailand reported 27.9% of cases with EONI is due to PROM.¹⁵ For such patients rapid diagnosis of early onset sepsis is needed, to avoid unnecessary stay in NICU, which in turn increase the economic and social burden in already poor settings. In agreement with the study done by Hornik¹⁶ and Altunhan¹⁷, CBC findings in this study was not informative, and didn't help in the diagnosis of sepsis. Also the British evidence update advisory group didn't recommend the use of CBC in the diagnosis of neonatal sepsis.¹⁸ Camacho A,¹⁹ stated that complete blood count is difficult to interpret in the neonatal period because it varies significantly with day of life and gestational age, and they are poor indicator of sepsis.

In this study the predominate organism isolated from patients with proven sepsis was Klebsiela 13%, followed by Ecoli 9%, this is in congruous with Zaidi et al.²⁰ and Downie L,²¹ who stated that Klebsiela is the most predominate organism in developing countries in both Hospital and community- acquired infection.

In contrast to other studies which stated that GBS is the most common pathogens of early onset sepsis in developed countries, but its burden in developing countries is less clear due to lack of studies using optimal diagnostic tools.²² GBS present in 8% of the patients of this study, done in a developing country. This can be explained, as the most frequent risk factor detected in this study was premature rupture of membrane and maternal infection which in turn increase the possibility of neonatal infection with GBS.

MRSA was detected in 2% of the patients in this study, nasal swabs from the NICU staff and swabs from the incubators were taken to detect the source of infection. Two nurses from the staff were MRSA positive; they were isolated and received treatment for 10 days.

Zaidi et al.²³ reported most pathogens isolated in the hospital setting before 72HR of life are similar to those isolated afterwards; it is likely that highly unclean delivery practices lead to infections with nosocomial agents very early in life.

Table 2 and 3 compare the results of CRP1 and PCT1 in the category of proven sepsis and not-proven sepsis. In both categories there was no significant difference between CRP1 and PCT1 in the ability to support the diagnosis of neonatal infection or refute it.

In the group with suspected sepsis there was no significant difference between PCT1 and CRP1 in distinguishing patients with neonatal sepsis.

Blommendahl J²⁴ stated that PCT was not a better marker than CRP levels because PCT is affected by perinatal factors within 48h of birth making its usefulness in diagnosing of early onset sepsis very limited. Other than infection, PCT levels increase in premature infants,

hypoxia, RDS, and hemodynamic instability, decreasing its specificity in early onset sepsis.²⁵ In this study CRP was repeated over 24 hours, while PCT was repeated after 8 hours to increase their reliability, the same was done in the study of Blommendahl et al,²⁴ and the study done by Hengest 2003.²⁶ Analysis of table 4 revealed that both CRP2 and PCT2 tests were able to differentiate more between infected and not infected newborns but there was no-significant difference between the two tests (P value 0.070). For the group of newborns with not-proven sepsis as shown in table 5, there was significant difference between both tests (P value 0.003), PCT2 CAN distinguish more accurately between cases with possible sepsis and cases with no sepsis.

Compares predictive values for CRP2 and PCT2 Sensitivity= 92.5%, Specificity=53%, PPV= 65.3%, NPV= 86.5%. PCT2: Sensitivity=95.8%, Specificity=87.7%, PPV=90.1%, NPV=98%. Procalcitonin evaluation done by Chaurasiya et al,²⁷ demonstrated sensitivity of 96.25%, specificity of 85%, PPV of 96.25% and NPV of 85%. Claudio Chiesa et al²⁸ studied the reliability of PCT concentration in 28 infants with severe early onset sepsis. They observed that the sensitivity 92.6%, specificity 97.5%, PPV 94.3%, NPV 96.5% respectively. It is clear from the previous data, that PCT didn't give additional information superior to that of CRP in diagnosing cases with suspected sepsis, in cases where blood culture results were not yet available. However, repeated PCT was more accurate than CRP in differentiating cases with sepsis. This means that the protocol of repeat test of PCT after 8 hours from the initial assessment could be of great benefit in diagnosing cases with early onset sepsis, thus reducing unnecessary stay in NICU, and over-use of antibiotics.

WHO developed the ASSURED criteria for an ideal point of care test in resource-limited settings, taking its accuracy and reliability as the main features.²⁹ PCT fulfills these criteria and the cost of assessment of PCT level twice for each patient is less than the cost of one day admission in NICU.

Making use of accuracy of PCT2 in detecting cases with neonatal sepsis, could have saved about 31 % of total admission days. Comparing this to the cost of the test, the benefit of assessing PCT is clear.

Conclusion

The expense of testing PCT twice is less than the expenses of one-day admission in NICU in developing countries. Application of such protocol could be of use in limiting period of stay in NICU.

References

1. Liu L, Johnson HL, Cousens S, Perin J, Scott S, Lawn JE et al. Global, regional, and national causes of child mortality: an updated systematic analysis for 2010 with time trends since 2000. *Lancet*. 2012; 379 (9832): 2151-61. doi: 10.1016/S0140-6736(12)60560-1.
2. National Neonatal Perinatal Database. Report 2002- 03. Published by NNPD Nodal Centre, Department of Pediatrics, All India Institute of Medical Science, New Delhi. Available from http://www.newbornwhocc.org/pdf/nnpd_report_2002-03. PDF last accessed 22 February 2019.
3. Vergnano S, Sharland M, Kazembe P, et al. Neonatal sepsis: an international perspective. *Arch Dis Child Fetal Neonatal Ed*. 2005 May; 90(3): F220-4. DOI:10. 1136/adc.2002.022863
4. Da Silva O, Ohlsson A, Kenyon C. Accuracy of leukocyte indices and C-reactive protein for diagnosis of neonatal sepsis: a critical review. *Pediatr Infect Dis J*. 1995 May;14(5):362-6.

5. Camacho-Gonzalez A, Spearman PW, Stoll BJ. Neonatal infectious diseases: evaluation of neonatal sepsis. *Pediatr Clin North Am.* 2013 Apr;60(2): 367- 89. doi: 10.1016/j.pcl.2012.12.003. Epub 2013 Jan 17.
6. Squire E, Favara B, Todd J. Diagnosis of neonatal bacterial infection: hematologic and pathologic findings in fatal and nonfatal cases. *Pediatrics.* 1979 Jul; 64 (1): 60-4.
7. Chauhan SB, Vaghasia V, Chauhan BB. C-reactive protein (crp) in early diagnosis of neonatal septicemia. *National Journal of Medical Research* 2012; 2 (3): 276–78.
8. Lakhey, A., & Shakya, H. Role of Sepsis Screening in early diagnosis of Neonatal Sepsis. *Journal of Pathology of Nepal*, 2017;7(1):1103-10. [https://doi.org /10.3126/jpn.v7i1.16944](https://doi.org/10.3126/jpn.v7i1.16944).
9. Pontrelli G, De Crescenzo F, Buzzetti R, Jenkner A, Balduzzi S, et al. Accuracy of serum procalcitonin for the diagnosis of sepsis in neonates and children with systemic inflammatory syndrome: A meta-analysis. *BMC Infect Dis.*2017;17(1):302. doi: 10.1186/s12879-017-2396-7.
10. Haque KN. Definitions of bloodstream infection in the newborn. *Pediatr Crit Care Med* 2005; 6(3 SUPPL):S45-9.
11. Mamta J., Kapil K., Garg LK., Vikas M., Mittal SK. To study the incidence and risk factors of early onset neonatal sepsis in an out born neonatal intensive care unit of India. *Clinical Neonatology* 2015, VOLUME 4, 91-95.
12. Khatua SP, Das AK, Chatterjee BD, Ghose B, Saha A. Neonatal septicemia. *Indian J Pediatr* 1986;53:509-14.
13. Muhammad Matloob Alam, Ali Faisal Saleem, Abdul Sattar Shaikah, Owais Munir, Maqbool Qadir. Neonatal sepsis following prolonged rupture of membranes in a tertiary care hospital in Karachi, Pakistan. *J Infect Dev Ctries* 2014; 8(1): 067-073.
14. Hossain MM AS, Shirin M, Chowdhury NA, Saha SK (2004). Bacterial etiology of neonatal sepsis in a tertiary care hospital in Bangladesh. *Bang J Child Health* 28:81-85.
15. Ratanakorn W, Srijarinya W, Chamnanvanakij S, Saengaroon P. (2005) Incidence of neonatal infection in newborn infants with a maternal history of premature rupture of membranes for 18 hours or longer by using phramongkutklao Hospital Clinical Practice Guidelines (CPG). *J Med Assoc Thai* 88:973-978.
16. Hornik CP, Benjamin DK, Becker KC, et al. Use of the complete blood count in early-onset neonatal sepsis. *Pediatr Infect Dis J* 2012; 31: 799-802.
17. Altunhan H, Annagur A, Ors, R., Mehmetoglu, I. Procalcitonin measurement at 24 hours of age may be helpful in the prompt diagnosis of early onset neonatal sepsis. *International Journal of Infectious Disease*, 15, E854-858, 2011
18. Mirrett, S., Weinstein, M.P., Reimer, L.G., Wilson, M. L., & Reller, L. (2001). Relevance of the number of positive bottles in determining clinical significance of coagulase-negative staphylococci in blood culture. *Journal of Clinical Microbiology*, 39, 3279-3281.
19. Camacho-Gonzalez A, Spearman PW, Stoll BJ. Neonatal infectious diseases: evaluation of neonatal sepsis. *Pediatr Clin North Am* 2013; 60: 367-89.
20. Zaidi AKM, Thaver D, Ali SA, et al. Pathogens associated with sepsis in newborns and young infants in developing countries. *Pediatr Infect Dis J* 2009; 28(1 Suppl): S10-81.
21. Downie L, Armiento R, Subhi R, et al. Community acquired neonatal and infant sepsis in developing countries: efficacy of WHO's currently recommended antibiotics-systemic review and meta-analysis. *Arch Dis Child* 2013; 98:146-54.
22. Pui-Ying Iroh Tam, Sherley F Delair, Stephen K Obaro. Neonatal group B streptococcus disease in developing countries. *Expert Review of vaccines*, volume 14, 1401-1403, 2015.

23. Zaidi AKM, Huskins WC, Thaver D, et al. Hospital acquired neonatal infections in developing countries. *Lancet* 2005; 365: 1175-88.
24. Blommendahl J, Janas M, Laine S, Miettinen A, Ashorn P. Comparison of procalcitonin with CRP and differential white blood cell count for the diagnosis of culture proven neonatal sepsis. *Scand J Infect Dis* 2002; 34: 620-22
25. Lapillonne A, Basson E, Monneret G, Bienvenu J, Salle BL. Lack of specificity of procalcitonin for sepsis diagnosis in premature infants. *Lancet* . 1998; 351:1211-12.
26. Hengest JM. The role of C-reactive protein in the evaluation and management of infants with suspected sepsis. *Advances in Neonatal Care*. 2003; 3:3-13.
27. Chaurasiya OS, Ahmed T, Chhabra K, Nath D, Hema J. Procalcitonin level in neonatal sepsis. *People's J of Scientific Research*. 2017; 10(1)
28. Claudio Chiesa, Pellegrini G, Alessandra Panero et al. C-RP, IL-6 and Procalcitonin in the immediate postnatal period: influence of illness severity, risk status, antenatal and perinatal complications and infection. *Clin Chem* 2003; 49(1):60-68.
29. Mabey D, Peeling RW, Ustianowski A, Perkins MD. Diagnostics for the developing world. *Nature Reviews Microbiology*. 2004. 2: 231-40.

Received: 16-09-2020 || Revised: 12-10-2020 || Accepted: 18-11-2020