

Original research article

## To Study the Menstrual Pattern in Cases of Women with Ovarian Masses: A Prospective Observational Research

Dr. Kiran Kumari

Senior Resident, Department of Obstetrics and Gynecology, Vardhman Institute of Medical Sciences, Pawapuri, Nalanda, Bihar, India

Corresponding Author: Dr. Kiran Kumari

### Abstract

**Aims:** To evaluate the menstrual characteristics in relation to ovarian cancer risk.

**Methods:** A prospective observational study was conducted in the Department of Obstetrics and Gynecology, Vardhman Institute of Medical Sciences, Pawapuri, Nalanda, Bihar, India, for 12 months. In all about 100 patients with palpable abdominal and pelvic adnexal mass with ovarian pathology were included for this study.

**Results:** Out of 100 patients 41% was 20-30 years age group and followed by 30-40 years. It is evident from the above table that 93 patients were married giving the incidence of 93% while 7% of cases were unmarried. 65% had regular menstrual cycle, 6% had surgical menopause, 3% had lactational amenorrhoea, 7% had amenorrhoea of pregnancy and 11% had physiological menopause, Secondary amenorrhea 2% and Bleeding P/V after amenorrhea was 2%.

**Conclusion:** There was no significant relation between menstrual history and occurrence of ovarian tumour.

**Keywords:** Menstrual, Woman, Ovarian & Masses.

### Introduction

Ovarian cancer is currently one of the most common forms of cancer among women worldwide, responsible for 3.6 % of all cases, with a mortality of 4.3 %. In Europe, it is the leading cause of cancer death among gynaecologic malignancies, ranking fifth in incidence (exceeded only by breast, colorectum, lung and corpus uteri) and sixth in mortality among all women's cancer (exceeded by breast, colorectum, lung, pancreas and stomach).<sup>1</sup> An important reason for the high mortality rates of this cancer is the late diagnosis. Many patients present in advanced stage, mostly because the disease is often asymptomatic or associated with nonspecific symptoms in the early stage. Incidental detection of adnexal mass is very common in clinical practice. Etiopathogenesis of ovarian tumours is not fully understood; however, it appears to be multifactorial. The leading risk factor is familiarity; namely, history of ovarian cancer in a first-degree relative. However, only 5–10 % of cases are related to hereditary syndromes: the main one is the breast-ovarian cancer syndrome, due to mutations in the BRCA1 and BRCA2 tumour suppressor genes. Approximately 90–95 % of cases are sporadic, with an increasing risk related to nulliparity, early menarche and late menopause, while pregnancy, lactation, early menopause and use of oral contraceptives appear to be protective factors.<sup>2</sup>

The influence of menstrual and reproductive factors, with the exception of parity, remains uncertain. The incessant ovulation hypothesis suggests that risk is increased by chronic post-ovulatory trauma to the epithelial surface of the ovary, and the tendency to form inclusion cysts.<sup>3</sup> The gonadotrophin hypothesis proposes that excessive gonadotropin secretion and consequent increases in oestrogen stimulation lead to proliferation and malignant

transformation of ovarian epithelium.<sup>4</sup> More recent hypotheses have suggested a role for chronic ovarian inflammation androgens and progesterone.<sup>5</sup> and the possibility that pregnancies reduce risk by clearing transforming cells from the ovaries.<sup>6</sup>

Up to 10% of women will have some form of surgery during their lifetime for the presence of an ovarian mass. In premenopausal women almost all ovarian masses and cysts are benign.<sup>7</sup> The overall incidence of a symptomatic ovarian cyst in a premenopausal female being malignant is approximately 1:1000 increasing to 3:1000 at the age of 50. Preoperative differentiation between the benign and the malignant ovarian mass in the premenopausal woman can be problematic with no test or algorithm being clearly superior in terms of accuracy.<sup>8</sup> Many ovarian masses in the premenopausal woman can be managed conservatively. Functional or simple ovarian cysts (thin-walled cysts without internal structures) which are less than 50 mm maximum diameter usually resolve over 2–3 menstrual cycles without the need for intervention.<sup>9</sup> we evaluate the menstrual characteristics in relation to ovarian cancer risk overall, and in relation to the major tumour histologic subtypes. Our findings are considered in light of several current hypotheses regarding ovarian cancer pathogenesis.

### Material and Methods

A prospective observational study was conducted in the Department of Obstetrics and Gynecology, Vardhman Institute of Medical Sciences, Pawapuri, Nalanda, Bihar, India, for 12 months. after taking the approval of the protocol review committee and institutional ethics committee.

### Methodology

All the patients with symptoms suggestive of adnexal mass were taken for the study and among these patients with ovarian pathology were subsequently included in the study. In all about 100 patients with palpable abdominal and pelvic adnexal mass with ovarian pathology were included for this study. The patients with uterine origin of mass were excluded from the study. A detailed history of each case was recorded with reference of age, religion, parity, socioeconomic status, symptomatology, marital status, menstrual history, obstetrics history, family history, history of contraceptive method, method adopted and history of present and past, medical and surgical illness. A special attention was given to those patients presenting with the four target symptoms viz. abdominal pain, abdominal mass, GIT symptoms and pelvic pain.

### Results

Out of 100 patients 41% was 20-30 years age group and followed by 30-40 years. It is evident from the above table that 93 patients were married giving the incidence of 93% while 7% of cases were unmarried. It is evident that 65% had regular menstrual cycle, 6% had surgical menopause, 3% had lactational amenorrhoea, 7% had amenorrhoea of pregnancy and 11% had physiological menopause, Secondary amenorrhea 2% and Bleeding P/V after amenorrhea was 2%. (Table 1-3)

**Table 1: Demographic profile of patients**

Parameter	N=100	%
Age		

Below 20 years	17	17
20-30 years	41	41
30-40 years	27	27
Above 40 years	15	15
<b>Education</b>		
Illiterate	17	17
Literate	83	83
<b>Occupation</b>		
House wife	79	79
Working	21	21
<b>Socio economic status</b>		
High	13	13
Middle	73	73
Low	14	14
<b>BMI</b>		
Normal	63	63
Over weight	37	37

**Table 2: Incidence of Ovarian Tumor According to Marital Status**

Marital Status	Ovarian tumours=100	Incidence
Married	93	93
Unmarried	7	7

**Table 3: Menstrual Pattern in Cases of Ovarian Tumors**

Menstrual Pattern	Number of Cases (N=100)	Incidence
Regular	65	65
Surgical menopause	6	6
Secondary amenorrhea	2	2
Bleeding P/V after amenorrhea	2	2
Lactational amenorrhea	3	3
Post-abortion amenorrhea	-	-
Amenorrhea of pregnancy	7	7
Physiological menopause	11	11
Oligomenorrhea	-	-
Menorrhagia	-	-
Polymenorrhagia	-	-
Continuous bleeding P/V	-	-
Metrorrhagia	-	-
Primary amenorrhea	-	-
Polymenorrhea	-	-

**Discussion**

Ovarian masses are a common finding in daily clinical practice and may be incidentally detected or identified in symptomatic patients. Characterization of an ovarian lesion represents

a diagnostic challenge; it is of great importance in the preoperative setting in order to plan adequate therapeutic procedures and may influence patient's management. The strong and consistently observed risk reduction associated with parity provides, in part, the basis of most hypotheses regarding ovarian pathogenesis. It has been shown, however, that interruption of ovulation during pregnancy, lactation, and oral contraceptive use is inadequate to account for the magnitude of the observed decrease in ovarian cancer risk.<sup>10</sup> Menstrual cycles occurring between ages 25 and 39 are most likely to be ovulatory<sup>11</sup> and pregnancies occurring between these ages have a greater potential to interrupt ovulatory cycles. Thus, our observation of reduced risk associated with later ages at first or last birth provides some support for hypotheses regarding incessant ovulation or ovarian inflammation. Pituitary secretion of gonadotropins generally increases during adulthood, but decreases during pregnancy; thus, the protective effects of later childbirth are also consistent with the gonadotropin hypothesis. The decreasing risk associated with later age at last birth is also consistent with the ovarian clearance hypothesis, and in particular with the notion that the protective 'clearance' effect of pregnancy is greater in older women, who are more likely to have premalignant ovarian epithelial cells.<sup>6</sup> Jeffcott's Principles of Gynaecology says that "Neither malignant nor benign growth usually affects the menstrual function in any way unless they happen to have sex endocrine function." Even if both ovaries are the seat of large tumours, there is always enough normal ovarian tissue left to continue a regular menstrual cycle.<sup>12</sup> In our study 65% had regular menstrual cycle, 6% had surgical menopause, 3% had lactational amenorrhoea, 7% had amenorrhoea of pregnancy and 11% had physiological menopause, Secondary amenorrhea 2% and Bleeding P/V after amenorrhea was 2%. Similar finding was observed other study.<sup>13</sup> Majority of the patients underwent surgical treatment excluding very few cases which were treated conservatively or were referred to cancer hospital.<sup>14</sup> 33(33%) patients underwent hysterectomy (TAH) among which 12(12%) underwent bilateral salpingo-oophorectomy, 7(7%) underwent unilateral salpingo-oophorectomy and 18(18%) with cyst removal. Ovarian cystectomy was done in 30(30%) cases. similar results was found in other study.<sup>15</sup> 5 patients were pregnant. In 5 cases LSCS was done and in 4 cases only the ovarian mass was removed in second trimester and the pregnancy continued. Mehta (1977)<sup>16</sup> reported incidence of ovariectomy and ovarian cystectomy to be 27.49% cases. Debulking surgery was done in 6% cases in my cases. Debulking was done due to advanced stage of the disease. All the patients were referred to cancer hospital for further management.

## Conclusion

The present study concluded that there was no significant relation between menstrual history and occurrence of ovarian tumour. Majority had regular menstrual cycle, 6% had surgical menopause, 3% had lactational amenorrhoea and Bleeding P/V after amenorrhea was 2%.

## Reference

1. Ferlay J, Soerjomataram I, Ervik M, Dikshit R, Eser S, Mathers C, Rebelo M, Parkin DM, Forman D, Bray F (2013) GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC Cancer Base No. 11. Lyon, France: International Agency for Research on Cancer. Available via <http://globocan.iarc.fr>, accessed on 01/03/2015 2.
2. Holschneider CH, Berek JS. Ovarian cancer: epidemiology, biology, and prognostic factors. *Semin Surg Oncol*.2000; 19(1):3–10
3. Casagrande JT, Louie EW, Pike MC, Roy S, Ross RK and Henderson BE. "Incessant ovulation" and ovarian cancer, *Lancet*.1979; 2: 170–173
4. Cramer DW and Welch WR . Determinants of ovarian cancer risk. II, Inferences regarding pathogenesis. *J Natl Cancer Inst*.1983; 71: 717–721

5. Risch HA. Hormonal etiology of epithelial ovarian cancer, with a hypothesis concerning the role of androgens and progesterone. *J Natl Cancer Inst.*1998; 90: 1774–1786
6. Adami H, Hsieh C, Lambe M, Trichopoulos D, Leon D and Persson I. Parity, age at first childbirth, and the risk of ovarian cancer. *Lancet.*1994; 344: 1250–1254
7. Lloyd H Smith, Cyllene R Morris, et al. Ovarian cancer: Can we make the clinical diagnosis earlier? *American Cancer Society.* 2005; 104:1398-407.
8. Singh Uma, Kohli Neera, Nisha, Ekta: Evaluation of new scoring system to differentiate between benign and malignant adnexal mass. *The Journal of Obstetrics and Gynecology of India.* 2006; 56(2):162-165.
9. Neerja Bhatla: Tumours of the ovary. In: Neerja Bhatla (Ed.) In: Jeffcoate's Principles of Gynaecology. London: Arnold, 5th Ed., 2001, 503-539
10. Risch JA, Weiss NS, Lyon JL, Daling JR and Liff JM. Events of reproductive life and the incidence of epithelial ovarian cancer. *Am J Epidemiol.*1983; 117: 128–139
11. Harlow SD and Ephross SA. Epidemiology of menstruation and its relevance to women's health. *Epidemiol Rev.*1995; 17: 265–286
12. Hans Olov Adami, David Hunter, Dimtrios Trichopoulos: Ovarian Cancer. In: Hans Olov Adami, David Hunter, Dimtrios Trichopoulos (Eds.): *Textbook of Cancer Epidemiology*, Oxford, 2002, 388.
13. Hans Olov Adami, David Hunter, Dimtrios Trichopoulos: Ovarian Cancer. In: Hans Olov Adami, David Hunter, Dimtrios Trichopoulos (Eds.): *Textbook of Cancer Epidemiology*, Oxford, 2002, 387.
14. David M Gerschem MD: Update on Epithelial Ovarian Cancer. In: David M. Gerschem (Ed.): *Obstetrics and Gynecology Clinics of North America*, 1994.
15. Lerner JP, Timor Tritsch IE, Federman A, Abramovich G: Transvaginal ultrasonographic characterization of ovarian masses with an improved weighted scoring system. *Am J Obstet Gynecol.* 1994; 170:81-5.
16. Mehta Buna: Study of cases of ovarian tumour. S.S. Medical College, Rewa, 1977

Received: 16-03-2021 // Revised: 13-04-2021 // Accepted: 03-05-2021