

Original research article

STUDY OF IRON DEFICIENCY ANEMIA ON GLYCATED HEMOGLOBIN (HBA1C) IN NON DIABETIC PATIENTS**Akanksha Srivastava¹, Ramesh Kumar², Niraj Diwakar³,
Sintu Kumar⁴, Sudhir Kumar⁵, Amit Kumar Mishra⁶**¹Junior Resident, Department of General medicine IGIMS Patna²Junior Resident, Department of General medicine IGIMS Patna³Junior Resident, Department of General medicine IGIMS Patna⁴Junior Resident, Department of General medicine IGIMS Patna⁵Prof. Department of General medicine IGIMS Patna⁶Additional Prof. Department of General medicine IGIMS Patna**Corresponding Author: Sintu Kumar****Abstract**

Background: The Iron deficiency anemia is the most common form of anemia in India. Hemoglobin A1c (HbA1c) is used in diabetic patients as an index of glycemic control reflecting glucose levels of the previous 3 months. Like blood sugar levels, HbA1c levels are also affected by the presence of variant hemoglobins, hemolytic anemias, nutritional anemias, uremia, pregnancy, and acute blood loss. We conducted a study to find out the influence of iron deficiency anaemia over HbAc levels.

Materials and Methods: A prospective observational study on 50 iron deficiency anaemia patient cases and 50 healthy control. Exclusion and inclusion criteria were used to recruit cases from the wards and OPDs of the hospital IN IGIMS Patna. HbA1c levels were measured at baseline and 3 months after treatment, and these values were compared with those in the control population.

Results: The HbA1C was significantly higher in the iron deficiency anaemic patients as compare to the control (5.88 ± 0.41 vs 5.03 ± 0.17 , respectively, $P < .05$). **Conclusion:** HbA1c is not affected by the blood sugar levels alone, and there various confounding factors when HbA1c is measured, especially that of iron deficiency, which is the commonest of the deficiency disease worldwide. rule out IDA before making a therapeutic decision, based on the HbA1c levels.

Keywords: Non-Diabetic Patient, Glycosylated Haemoglobin, IDA, HbA1c.

Introduction

Diabetes mellitus is a metabolic disease which is caused by absolute or relative insulin deficiency . about 10% of the Indian population suffers from this disease.HbA1c is used as the gold standard for monitoring glycaemic control and used as a foresee of diabetic complications.1 Throughout the circulatory life of the red blood cell, HbA1c is formed continuously by adjoining of glucose to the Nterminal of the haemoglobin beta chain. This process is non-enzymatic and reflects the average exposure of haemoglobin to glucose over a period of nearly 70 - 90 days. HbA1c has been defined as the fast fraction haemoglobin (HbA1a, A1c), which elute first during column chromatography with cation-exchange resin. HbA1c levels are not influenced by only blood glucose levels. They are also altered in

hemolytic anemias, hemoglobinopathies, acute and chronic blood loss, pregnancy, and uremia, Vitamin B12, folate, and iron deficiency anemias have also been shown to affect HbA1c levels². The two known factors which can modulate the glycation of proteins are the prevailing concentration of glucose and the half life of the protein³. But evidences in the literature have documented increased glycated protein levels in some nondiabetic pathological states, like iron deficiency anemia. Some authors have also found that on supplementation with iron therapy, there was a significant decrease in the levels of glycated haemoglobin⁴. Evidence has accumulated, which supports the hypothesis that the glycation reaction, apart from the traditional chronic hyperglycaemia, can be modulated by the iron status of the patient. If the degree of glycation of other proteins in anemic patients was similar to that of the glycated haemoglobin, it would have important clinical implications. Some studies show that HbA1c levels are increased in iron deficiency anemia and attempted to explain on the basis of both modifications to the structure of haemoglobin and level of HbA1c in old and new red blood cells⁵. Some Study showed that HbA1c levels were higher in patients of iron deficiency anemia and decreased significantly upon treatment with iron⁶. With this background, the objective of the present study was to study the effect of iron deficiency anemia on glycated hemoglobin (HbA1c) in non diabetic Indian subjects. The anaemic patients were selected, based on their haemoglobin levels (Hb < 11g/dl), and on their peripheral blood smears (mostly microcytic hypochromic), which suggested iron deficiency anaemia and on their haematologic investigations (MCV < MCH < MCHC) and serum fasting and postprandial glucose levels.

MATERIALS AND METHODS

Type of Study- Prospective observational study conducted in Indira Gandhi Institute of Medical Sciences patna bihar. The iron deficiency anaemic patients were recruited on their haemoglobin levels (Hb < 12 in non-pregnant female and 13 in male), ferritin levels (< 9 ng/mL for women, < 15 ng/mL for men) and on their peripheral blood smears which indicate iron deficiency anaemia. The levels of haemoglobin, mean corpuscular haemoglobin (MCH), haematocrit, mean corpuscular volume (MCV), mean corpuscular haemoglobin concentration (MCHC), platelet count, total leucocyte count (TLC) and differential leucocyte count (DLC) were measured by an automated analyser. On the basis of haemoglobin level patients were categorised as having mild, moderate or severe anaemia. Mild anaemia (male patients 12 - 12.9 gm/ dL and female patients 11 - 11.9 gm/ dL). Moderate anaemia (male patients 9 - 11.9 gm/ dL and female patients 8 - 10.9 gm/ dL) and severe anaemia (male patients < 9 gm/ dL and female patients < 8 gm/ dL). Those with predominantly microcytic indices (MCV < 80 fL), hypochromic indices (MCH < 26 pg/ cell) were considered to have iron deficiency anaemia. Normal HbA1c in nondiabetic individuals is < 5.6%.

Inclusion Criteria

1. Patient without diabetes.
2. Patient belonging to rural and urban area.
3. > 18 yrs. of age.
4. Giving consent for study.

Exclusion Criteria

Patients with history of acute or chronic blood loss, haemolytic anaemia, haemoglobinopathies, kidney diseases, diabetics, pregnancy, chronic alcohol ingestion and impaired glucose tolerance are excluded. The blood specimens were drawn after an 8-

hourly fasting period. A Sysmex automated haematology analyser (MINDRAY) was used for the whole blood counts [haemoglobin (Hb), mean corpuscular volume (MCV) and mean corpuscular haemoglobin (MCH)]. The serum ferritin levels were measured by using a Diatek kit in a Labowind semiautomated analyser and the peripheral blood smears were examined in all the patients. The HbA1c levels were determined by turbidimetric immunoinhibition.

RESULTS:

The present study was conducted among 50 cases having Iron deficiency anemia and 50 apparently healthy controls without iron deficiency anemia. All cases and controls are non diabetic. all the oarameters which were tested in both the groups,the fasting and the postprandial blood glucose levels confirmed the non diabetic status.

Table 1: Demographic variable

Variable	Case group	Control group	p-value
Age (Yrs)	48.23±4.6	47.23±4.92	>0.05
Male : Female	12:38	13:37	>0.05

There were no significant differences between the two groups with regard to age and gender.

Table 2: Comparison of blood parameter

Variable	Case group	Control group	p-value
HB (Gm/dl)	6.01±1.9	12.4±1.2	<0.001
S. Ferritin (ng/mL)	8.71±1.68	90.21±12.54	<0.001
HbA1C	5.88±0.41	5.03±0.17	<0.05

The HbA1C was significantly higher in the iron deficiency patients as compare to the control (5.88 ± 0.41 vs 5.03 ± 0.17, respectively, P < .05). **DISCUSSION** Our results suggested that IDA was associated with higher concentrations of HbA1c. The mean HbA1c of cases was 5.88% while it was 5.03% in healthy control. Application of statistical test shows that the difference of HbA1c between case and control group was statistically significant. In the study conducted by Vishal Kalasker et al , the mean HbA1c of cases was 5.91±0.47 % while it was 6.54±0.39 % in healthy control.

Application of statistical test shows that the difference of HbA1c between case and control group was statistically significant indicating statistically higher HbA1c in control group. 7 In the study conducted by Balasubramanian Shanthi et al , the mean HbA1c of cases was 7.6 ± 0.5 % while it was 5.5 ± 0.8 % in healthy control. Application of statistical test shows that the difference of HbA1c between case and control group was statistically significant indicating statistically higher HbA1c in cases group. 8 In the study conducted by Alap L. Christy et al , the mean HbA1c of cases was 6.87 ± 1.4% while it was 5.65 ± 0.69% in healthy control. Application of statistical test shows that the difference of HbA1c, iron deficiency anaemia is most common nutritional anaemia, HbA1c was used not only to asses the glycaemic control but also for diagnosis and screening for diabetes. between case and control group was statistically significant indicating statistically higher HbA1c in cases group.8

CONCLUSION

Our study we would like to conclude that iron deficiency was associated with higher proportions of HbA1c, which could cause problems in the diagnosis of uncontrolled diabetes mellitus in iron-deficient patients.

REFERENCES

1. Telen MJ, Kaufman RE. The mature erythrocyte. In: Greer JP, Forester J, et al., editors. *Wintrobe's clinical hematology*. 11th ed. Lippincot: Williams and Wilkins; 2004. p. 230
2. American Diabetes Association. Standards of medical care in diabetes—2016. *Diabetes Care*. 2016;39(suppl 1):S1- S106.
3. Starkman HS, wacks M, Soeldner JS, Kim A. Effect of acute blood loss on glycosylated hemoglobin determination in normal subjects. *Diabetes care* 1983;6:291-4
4. Mayer TK, Freedman ZR. Protein glycosylation in diabetes mellitus: a review of the laboratory measurements and their clinical utilities. *Clin Chim Acta*. 1983;127: 147e 84
5. Chowdhury TA, Lasker SS. The elevated glycated haemoglobin in nondiabetic patients is associated with an increased mortality in myocardial infarction. *Postgrad Med J*. 1998;74:480e1
6. Braunwald E, Fauci AS, Kasper DL, Hauser SL, Longo DL, Jameson JL: Diabetes mellitus. *Harrison's Principles of Internal Medicine*, ed 17, 2008; 2275-23
7. Vishal Kalasker , Sudhamadhuri, M V Kodliwadmath, Harish Bhat. Effect of Iron Deficiency Anemia on Glycosylated Hemoglobin Levels in Non Diabetic Indian Adults. *International Journal of Medical and Health Sciences*, Jan 2014; Vol-3; Issue-1 p-40-43.
8. Balasubramanian Shanthi et al., Effect of Iron Deficiency on Glycation of Hemoglobin *Journal of Clinical and Diagnostic Research*. 2013 January, Vol- 7(1): 15-17
9. Alap L. Christy, Poornima A. Manjrekar*, Ruby P. Babu, Anupama Hegde, and Rukmini M.S. Influence of Iron Deficiency Anemia on Hemoglobin A1C Levels in Diabetic Individuals with Controlled Plasma Glucose Levels. *Iranian Biomedical Journal*, April 2014; 18 (2): 88-93
10. Van Heyningen C, Dalton RG, Glycosylated haemoglobin in iron deficiency anaemia. *Lancet*. 1985; 1: 874.
11. Rai KB, Pattaabiraman TN. Glycosylated haemoglobin levels in iron deficiency anaemia. *Indian J Med Res*. 1980; 83: 234-6